

CS434a/541a: Pattern Recognition
Prof. Olga Veksler

Lecture 6

Today

- Introduction to nonparametric techniques
- Basic Issues in Density Estimation
- Two Density Estimation Methods
 1. Parzen Windows
 2. Nearest Neighbors

Non-Parametric Methods

- Neither probability distribution nor discriminant function is known
 - Happens quite often
- All we have is labeled data



- Estimate the probability distribution from the labeled data

*a lot is known
"easier"*



*little is known
"harder"*

NonParametric Techniques: Introduction

- In previous lectures we assumed that either
 1. someone gives us the density $p(\mathbf{x})$
 - In pattern recognition applications this never happens
 2. someone gives us $p(\mathbf{x}/\theta)$
 - Does happen sometimes, **but**
 - we are likely to suspect whether the given $p(\mathbf{x}/\theta)$ models the data well
 - Most parametric densities are unimodal (have a single local maximum), whereas many practical problems involve multi-modal densities

NonParametric Techniques: Introduction


- Nonparametric procedures can be used with arbitrary distributions and without any assumption about the forms of the underlying densities
- There are two types of nonparametric methods:
 - Parzen windows
 - Estimate likelihood $p(\mathbf{x} / \mathbf{c}_j)$
 - Nearest Neighbors
 - Bypass likelihood and go directly to posterior estimation $P(\mathbf{c}_j / \mathbf{x})$

NonParametric Techniques: Introduction

- Nonparametric techniques attempt to estimate the underlying density functions from the training data
 - Idea: the more data in a region, the larger is the density function

$$Pr[X \in \mathcal{R}] = \int_{\mathcal{R}} f(x) dx$$

*average of $f(x)$
over \mathcal{R}*



NonParametric Techniques: Introduction

$$Pr[X \in \mathcal{R}] = \int_{\mathcal{R}} f(x) dx$$

- How can we approximate $Pr[X \in \mathcal{R}_1]$ and $Pr[X \in \mathcal{R}_2]$?


- $Pr[X \in \mathcal{R}_1] \approx \frac{6}{20}$ and $Pr[X \in \mathcal{R}_2] \approx \frac{6}{20}$

- Should the density curves above \mathcal{R}_1 and \mathcal{R}_2 be equally high?

- No, since \mathcal{R}_1 is smaller than \mathcal{R}_2

$$Pr[X \in \mathcal{R}_1] = \int_{\mathcal{R}_1} f(x) dx \approx \int_{\mathcal{R}_2} f(x) dx = Pr[X \in \mathcal{R}_2]$$

- To get density, normalize by region size



NonParametric Techniques: Introduction

- Assuming $f(\mathbf{x})$ is basically flat inside \mathcal{R} ,

$$\frac{\text{\# of samples in } \mathcal{R}}{\text{total \# of samples}} \approx \text{Pr}[X \in \mathcal{R}] = \int_{\mathcal{R}} f(\mathbf{y}) d\mathbf{y} \approx f(\mathbf{x}) * \text{Volume}(\mathcal{R})$$

- Thus, density at a point \mathbf{x} inside \mathcal{R} , can be approximated

$$f(\mathbf{x}) \approx \frac{\text{\# of samples in } \mathcal{R}}{\text{total \# of samples}} \frac{1}{\text{Volume}(\mathcal{R})}$$

- Now let's derive this formula more formally

Binomial Random Variable

- Let us flip a coin n times (each one is called “trial”)
 - Probability of head ρ , probability of tail is $1-\rho$
- Binomial random variable K counts the number of heads in n trials

$$P(K = k) = \binom{n}{k} \rho^k (1 - \rho)^{n-k}$$

where $\binom{n}{k} = \frac{n!}{k!(n-k)!}$

- Mean is $E(K) = n\rho$
- Variance is $\text{var}(K) = n\rho(1 - \rho)$

Density Estimation: Basic Issues

- From the definition of a density function, probability ρ that a vector \mathbf{x} will fall in region \mathcal{R} is:

$$\rho = \Pr[\mathbf{x} \in \mathcal{R}] = \int_{\mathcal{R}} p(\mathbf{x}') d\mathbf{x}'$$

- Suppose we have samples $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ drawn from the distribution $p(\mathbf{x})$. The probability that k points fall in \mathcal{R} is then given by binomial distribution:

$$\Pr[K = k] = \binom{n}{k} \rho^k (1 - \rho)^{n-k}$$

- Suppose that k points fall in \mathcal{R} , we can use MLE to estimate the value of ρ . The likelihood function is


$$p(\mathbf{x}_1, \dots, \mathbf{x}_n | \rho) = \binom{n}{k} \rho^k (1 - \rho)^{n-k}$$

Density Estimation: Basic Issues

$$p(\mathbf{x}_1, \dots, \mathbf{x}_n | \rho) = \binom{n}{k} \rho^k (1 - \rho)^{n-k}$$

- This likelihood function is maximized at $\rho = \frac{k}{n}$
- Thus the MLE is $\hat{\rho} = \frac{k}{n}$
- Assume that $p(\mathbf{x})$ is continuous and that the region \mathcal{R} is so small that $p(\mathbf{x})$ is approximately constant in \mathcal{R}

$$\int_{\mathcal{R}} p(\mathbf{x}') d\mathbf{x}' \cong p(\mathbf{x})V$$




- \mathbf{x} is in \mathcal{R} and V is the volume of \mathcal{R}
- Recall from the previous slide: $\rho = \int_{\mathcal{R}} p(\mathbf{x}') d\mathbf{x}'$
- Thus $p(\mathbf{x})$ can be approximated: $p(\mathbf{x}) \approx \frac{k/n}{V}$

Density Estimation: Basic Issues

- This is exactly what we had before:

$$p(\mathbf{x}) \approx \frac{k/n}{V}$$



x is inside some region \mathcal{R}

k = number of samples inside \mathcal{R}

n = total number of samples inside \mathcal{R}

V = volume of \mathcal{R}


- Our estimate will always be the average of true density over \mathcal{R}

$$p(\mathbf{x}) \approx \frac{k/n}{V} = \frac{\hat{\rho}}{V} \approx \frac{\int_{\mathcal{R}} p(\mathbf{x}') d\mathbf{x}'}{V}$$

- Ideally, $p(\mathbf{x})$ should be constant inside \mathcal{R}

Density Estimation: Histogram


$$p(x) \approx \frac{k/n}{V}$$



- If regions \mathcal{R}_i 's do not overlap, we have a histogram

Density Estimation: Accuracy

- How accurate is density approximation $p(\mathbf{x}) \approx \frac{k/n}{V}$?
- We have made two approximations
 1. $\hat{\rho} = \frac{k}{n}$
 - as n increases, this estimate becomes more accurate
 2. $\int_{\mathcal{R}} p(\mathbf{x}') d\mathbf{x}' \cong p(\mathbf{x})V$
 - as \mathcal{R} grows smaller, the estimate becomes more accurate
 - As we shrink \mathcal{R} we have to make sure it contains samples, otherwise our estimated $p(\mathbf{x}) = 0$ for all \mathbf{x} in \mathcal{R}
- Thus in theory, if we have an unlimited number of samples, to we get convergence as we simultaneously increase the number of samples n , and shrink region \mathcal{R} , but not too much so that \mathcal{R} still contains a lot of samples



Density Estimation: Accuracy

$$p(\mathbf{x}) \approx \frac{k/n}{V}$$


- In practice, the number of samples is always fixed
- Thus the only available option to increase the accuracy is by decreasing the size of \mathcal{R} (V gets smaller)
 - If V is too small, $p(\mathbf{x})=0$ for most \mathbf{x} , because most regions will have no samples
 - Thus have to find a compromise for V
 - not too small so that it has enough samples
 - but also not too large so that $p(\mathbf{x})$ is approximately constant inside V

Density Estimation: Two Approaches

$$p(\mathbf{x}) \approx \frac{k/n}{V}$$


1. Parzen Windows:

- Choose a fixed value for volume V and determine the corresponding k from the data




2. k-Nearest Neighbors

- Choose a fixed value for k and determine the corresponding volume V from the data
- Under appropriate conditions and as number of samples goes to infinity, both methods can be shown to converge to the true $p(\mathbf{x})$




Parzen Windows


- In Parzen-window approach to estimate densities we fix the size and shape of region \mathcal{R}
- Let us assume that the region \mathcal{R} is a d -dimensional hypercube with side length h thus it's volume is h^d



1 dimension



2 dimensions




3 dimensions

Parzen Windows

- To estimate the density at point \mathbf{x} , simply center the region \mathcal{R} at \mathbf{x} , count the number of samples in \mathcal{R} , and substitute everything in our formula

$$p(\mathbf{x}) \approx \frac{k/n}{V}$$





$$p(\mathbf{x}) \approx \frac{3/6}{10}$$

Parzen Windows

- We wish to have an analytic expression for our approximate density \mathcal{R}
- Let us define a **window function**


$$\varphi(u) = \begin{cases} 1 & |u_j| \leq \frac{1}{2} \quad j = 1, \dots, d \\ 0 & \text{otherwise} \end{cases}$$



Parzen Windows

- Recall we have samples $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$. Then

$$\varphi\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right) = \begin{cases} 1 & |\mathbf{x} - \mathbf{x}_i| \leq \frac{h}{2} \quad j = 1, \dots, d \\ 0 & \text{otherwise} \end{cases}$$



$$\varphi\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right) = \begin{cases} 1 & \text{if } \mathbf{x}_i \text{ is inside the hypercube with} \\ & \text{width } h \text{ and centered at } \mathbf{x} \\ 0 & \text{otherwise} \end{cases}$$

Parzen Windows

- How do we count the total number of sample points $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ which are inside the hypercube with side h and centered at \mathbf{x} ?

$$k = \sum_{i=1}^{i=n} \varphi\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right)$$

- Recall $p(\mathbf{x}) \approx \frac{k/n}{V}$
- Thus we get the desired analytical expression for the estimate of density $p_\varphi(\mathbf{x})$

$$p_\varphi(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^{i=n} \frac{1}{h^d} \varphi\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right)$$

Parzen Windows

$$p_{\varphi}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^{i=n} \frac{1}{h^d} \varphi\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right)$$

- Let's make sure $p_{\varphi}(\mathbf{x})$ is in fact a density

- $p_{\varphi}(\mathbf{x}) \geq 0 \quad \forall \mathbf{x}$

volume of hypercube

- $$\int p_{\varphi}(\mathbf{x}) d\mathbf{x} = \int \frac{1}{n} \sum_{i=1}^{i=n} \frac{1}{h^d} \varphi\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right) d\mathbf{x} = \frac{1}{h^d n} \sum_{i=1}^{i=n} \int \varphi\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right) d\mathbf{x}$$
$$= \frac{1}{n} \frac{1}{h^d} \sum_{i=1}^{i=n} h^d = 1$$

Today

- Continue nonparametric techniques
 1. Finish Parzen Windows
 2. Start Nearest Neighbors (hopefully)

Parzen Windows

$$p(\mathbf{x}) \approx \frac{k/n}{V}$$


x is inside some region \mathcal{R}

k = number of samples inside \mathcal{R}

n = total number of samples inside \mathcal{R}

V = volume of \mathcal{R}

- To estimate the density at point \mathbf{x} , simply center the region \mathcal{R} at \mathbf{x} , count the number of samples in \mathcal{R} , and substitute everything in our formula



$$p(\mathbf{x}) \approx \frac{3/6}{10}$$

Parzen Windows

- Formula for Parzen window estimation


$$p_{\varphi}(\mathbf{x}) = \frac{\sum_{i=1}^n \varphi\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right)}{h^d} = \frac{1}{n} \sum_{i=1}^n \frac{1}{h^d} \varphi\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right)$$

The diagram includes handwritten annotations: a green oval around the summation term is labeled "= k", and a red oval around the h^d term in the denominator is labeled "= V".

Parzen Windows: Example in 1D

$$p_{\varphi}(x) = \frac{1}{n} \sum_{i=1}^n \frac{1}{h^d} \varphi\left(\frac{x - x_i}{h}\right)$$

- Suppose we have 7 samples $D = \{2, 3, 4, 8, 10, 11, 12\}$



- Let window width $h=3$, estimate density at $x=1$



$$p_{\varphi}(1) = \frac{1}{7} \sum_{i=1}^7 \frac{1}{3} \varphi\left(\frac{1 - x_i}{3}\right) = \frac{1}{21} \left[\varphi\left(\frac{1-2}{3}\right) + \varphi\left(\frac{1-3}{3}\right) + \varphi\left(\frac{1-4}{3}\right) + \dots + \varphi\left(\frac{1-12}{3}\right) \right]$$

$$\left| -\frac{1}{3} \right| \leq 1/2 \quad \left| -\frac{2}{3} \right| > 1/2 \quad \left| -1 \right| > 1/2 \quad \left| -\frac{11}{3} \right| > 1/2$$

$$p_{\varphi}(1) = \frac{1}{7} \sum_{i=1}^7 \frac{1}{3} \varphi\left(\frac{1 - x_i}{3}\right) = \frac{1}{21} [1 + 0 + 0 + \dots + 0] = \frac{1}{21}$$

Parzen Windows: Sum of Functions

- Fix \mathbf{x} , let i vary and ask
 - For which samples \mathbf{x}_i is $\varphi\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right) = 1$?



- Now fix \mathbf{f} and let \mathbf{x} vary and ask
 - For which \mathbf{x} is $\varphi\left(\frac{\mathbf{x} - \mathbf{x}_f}{h}\right) = 1$? For all \mathbf{x} in gray box
- Thus $\varphi\left(\frac{\mathbf{x} - \mathbf{x}_f}{h}\right) = 1$ is simply a function which is 1 inside square of width h centered at \mathbf{x}_f and 0 otherwise!

Parzen Windows: Sum of Functions

- Now let's look at our density estimate $\hat{p}_\varphi(\mathbf{x})$ again:


$$\hat{p}_\varphi(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^{i=n} \frac{1}{h^d} \varphi\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right) = \sum_{i=1}^{i=n} \underbrace{\frac{1}{nh^d} \varphi\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right)}$$

*1 inside square centered at \mathbf{x}_i
0 otherwise*

- Thus $\hat{p}_\varphi(\mathbf{x})$ is just a sum of n “box like” functions each of height $\frac{1}{nh^d}$

Parzen Windows: Example in 1D

- Let's come back to our example
 - 7 samples $D=\{2,3,4,8,10,11,12\}$, $h=3$




- To see what the function looks like, we need to generate 7 boxes and add them up
- The width is $h=3$ and the height, according to previous slide is

$$\frac{1}{nh^d} = \frac{1}{21}$$


Parzen Windows: Interpolation

- In essence, window function ϕ is used for interpolation: each sample \mathbf{x}_i contributes to the resulting density at \mathbf{x} if \mathbf{x} is close enough to \mathbf{x}_i




Parzen Windows: Drawbacks of Hypercube φ

- As long as sample point \mathbf{x}_i and \mathbf{x} are in the same hypercube, the contribution of \mathbf{x}_i to the density at \mathbf{x} is constant, regardless of how close \mathbf{x}_i is to \mathbf{x}



$$\varphi\left(\frac{\mathbf{x} - \mathbf{x}_1}{h}\right) = \varphi\left(\frac{\mathbf{x} - \mathbf{x}_2}{h}\right) = 1$$

- The resulting density $p_\varphi(\mathbf{x})$ is not smooth, it has discontinuities



Parzen Windows: general φ

$$p_{\varphi}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n \frac{1}{h^d} \varphi\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right)$$

- We can use a general window φ as long as the resulting $p_{\varphi}(\mathbf{x})$ is a legitimate density, i.e.

1. $p_{\varphi}(\mathbf{u}) \geq 0$


- satisfied if $\varphi(\mathbf{u}) \geq 0$

2. $\int p_{\varphi}(\mathbf{x}) d\mathbf{x} = 1$

- satisfied if $\int \varphi(\mathbf{u}) d\mathbf{u} = 1$

$$\int p_{\varphi}(\mathbf{x}) d\mathbf{x} = \frac{1}{nh^d} \sum_{i=1}^n \int \varphi\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right) d\mathbf{x} = \frac{1}{nh^d} \sum_{i=1}^n \int h^d \varphi(\mathbf{u}) d\mathbf{u} = 1$$


change coordinates to $u = \frac{\mathbf{x} - \mathbf{x}_i}{h}$, thus $du = \frac{d\mathbf{x}}{h}$



Parzen Windows: general φ

$$p_{\varphi}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n \frac{1}{h^d} \varphi\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right)$$

- Notice that with the general window φ we are no longer counting the number of samples inside \mathcal{R} .
- We are counting the weighted average of potentially every single sample point (although only those within distance h have any significant weight)



- With infinite number of samples, and appropriate conditions, it can still be shown that


$$p_{\varphi}^n(\mathbf{x}) \rightarrow p(\mathbf{x})$$

Parzen Windows: Gaussian φ

$$p_{\varphi}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n \frac{1}{h^d} \varphi\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right)$$

- A popular choice for φ is $\mathbf{N}(\mathbf{0}, \mathbf{1})$ density

$$\varphi(u) = \frac{1}{\sqrt{2\pi}} e^{-u^2/2}$$




- Solves both drawbacks of the “box” window
 - Points \mathbf{x} which are close to the sample point \mathbf{x}_i receive higher weight
 - Resulting density $p_{\varphi}(\mathbf{x})$ is smooth

Parzen Windows: Example with General φ

- Let's come back to our example
 - 7 samples $D=\{2,3,4,8,10,11,12\}$, $h=1$


$$p_{\varphi}(\mathbf{x}) = \frac{1}{7} \sum_{i=1}^{i=7} \varphi(\mathbf{x} - \mathbf{x}_i)$$




- $p_{\varphi}(\mathbf{x})$ is the sum of 7 Gaussians, each centered at one of the sample points, and each scaled by $1/7$

Parzen Windows: Did We Solve the Problem?

- Let's test if we solved the problem
 1. Draw samples from a known distribution
 2. Use our density approximation method and compare with the true density
- We will vary the number of samples n and the window size h
- We will play with 2 distributions




$N(0, 1)$



***triangle and
uniform mixture***

Parzen Windows: True Density $N(0,1)$



Parzen Windows: True Density $N(0,1)$





FIGURE 4.5. Parzen-window estimates of a univariate normal density using different window widths and numbers of samples. The vertical axes have been scaled to best show the structure in each graph. Note particularly that the $n = \infty$ estimates are the same (and match the true density function), regardless of window width. From: Richard

Parzen Windows: True density is Mixture of Uniform and Triangle



Parzen Windows: True density is Mixture of Uniform and Triangle

$h=1$

$h=0.5$

$h=0.2$





FIGURE 4.7. Parzen-window estimates of a bimodal distribution using different window widths and numbers of samples. Note particularly that the $n = \infty$ estimates are the same (and match the true distribution), regardless of window width. From: Richard O. Duda, Peter E. Hart, and David G. Stork, *Pattern Classification*. Copyright © 2001 by John Wiley & Sons, Inc.

Parzen Windows: Effect of Window Width h

- By choosing h we are guessing the region where density is approximately constant
- Without knowing anything about the distribution, it is really hard to guess where the density is approximately constant




Parzen Windows: Effect of Window Width h

- If h is small, we superimpose n sharp pulses centered at the data
 - Each sample point \mathbf{x}_i influences too small range of \mathbf{x}
 - **Smoothed too little**: the result will look noisy and not smooth enough
- If h is large, we superimpose broad slowly changing functions,
 - Each sample point \mathbf{x}_i influences too large range of \mathbf{x}
 - **Smoothed too much**: the result looks oversmoothed or “out-of-focus”
- Finding the best h is challenging, and indeed no single h may work well
 - May need to adapt h for different sample points
- **However we can try to learn the best h to use from the test data**


Parzen Windows: Classification Example

- In classifiers based on Parzen-window estimation:
 - We estimate the densities for each category and classify a test point by the label corresponding to the maximum posterior
 - The decision region for a Parzen-window classifier depends upon the choice of window function as illustrated in the following figure

Parzen Windows: Classification Example



- For **small** enough window size h classification on training data is be perfect
- However decision boundaries are complex and this solution is not likely to generalize well to novel data



- For **larger** window size h , classification on training data is not perfect
- However decision boundaries are simpler and this solution is more likely to generalize well to novel data

Parzen Windows: Summary

- Advantages
 - Can be applied to the data from any distribution
 - In theory can be shown to converge as the number of samples goes to infinity
- Disadvantages
 - Number of training data is limited in practice, and so choosing the appropriate window size h is difficult
 - May need large number of samples for accurate estimates
 - Computationally heavy, to classify one point we have to compute a function which potentially depends on all samples
 - Window size h is not trivial to choose

k-Nearest Neighbors


- Recall the generic expression for density estimation

$$p(\mathbf{x}) \approx \frac{k / n}{V}$$

- In Parzen windows estimation, we fix V and that determines k , the number of points inside V
- In k-nearest neighbor approach we fix k , and find V that contains k points inside

k-Nearest Neighbors

- kNN approach seems a good solution for the problem of the “best” window size
 - Let the cell volume be a function of the training data
 - Center a cell about x and let it grow until it captures k samples
 - k are called the k nearest-neighbors of x




- 2 possibilities can occur:
 - Density is high near x ; therefore the cell will be small which provides a good resolution
 - Density is low; therefore the cell will grow large and stop until higher density regions are reached

k-Nearest Neighbor

- Of course, now we have a new question
 - How to choose k ?
- A good “rule of thumb” is $k = \sqrt{n}$
 - Can prove convergence if n goes to infinity
 - Not too useful in practice, however
- Let’s look at 1-D example
 - we have one sample, i.e. $n = 1$


$$p(\mathbf{x}) \approx \frac{k/n}{V} = \frac{1}{2|\mathbf{x} - \mathbf{x}_1|}$$



- But the estimated $p(\mathbf{x})$ is not even close to a density function:

$$\int_{-\infty}^{\infty} \frac{1}{2|\mathbf{x} - \mathbf{x}_1|} d\mathbf{x} = \infty \neq 1$$

k-Nearest Neighbor: Gaussian and Uniform plus Triangle Mixture Estimation



k-Nearest Neighbor: Gaussian and Uniform plus Triangle Mixture Estimation




FIGURE 4.12. Several k -nearest-neighbor estimates of two unidimensional densities: a Gaussian and a bimodal distribution. Notice how the finite n estimates can be quite “spiky.” From: Richard O. Duda, Peter E. Hart, and David G. Stork, *Pattern Classification*. Copyright © 2001 by John Wiley & Sons, Inc.


Today

- Continue with Nonparametric Density Estimation
 - Finish Nearest Neighbor

k-Nearest Neighbors

$$p(\mathbf{x}) \approx \frac{k / n}{V}$$

- kNN approach seems a good solution for the problem of the “best” window size
 - Let the cell volume be a function of the training data
 - Center a cell about \mathbf{x} and let it grow until it captures k samples
 - k are called the k nearest-neighbors of \mathbf{x}



k-Nearest Neighbor: Gaussian and Uniform plus Triangle Mixture Estimation




FIGURE 4.12. Several k -nearest-neighbor estimates of two unidimensional densities: a Gaussian and a bimodal distribution. Notice how the finite n estimates can be quite “spiky.” From: Richard O. Duda, Peter E. Hart, and David G. Stork, *Pattern Classification*. Copyright © 2001 by John Wiley & Sons, Inc.

k-Nearest Neighbor

- Thus straightforward density estimation $p(\mathbf{x})$ does not work very well with kNN approach because the resulting density estimate
 1. Is not even a density
 2. Has a lot of discontinuities (looks very spiky, not differentiable)
 3. Even for large regions with no observed samples the estimated density is far from zero (tails are too heavy)
- *Notice in the theory, if infinite number of samples is available, we could construct a series of estimates that converge to the true density using kNN estimation. However this theorem is not very useful in practice because the number of samples is always limited*

k-Nearest Neighbor

- However we shouldn't give up the nearest neighbor approach yet
- Instead of approximating the density $p(\mathbf{x})$, we can use kNN method to approximate the posterior distribution $P(\mathbf{c}_i|\mathbf{x})$
 - We don't even need $p(\mathbf{x})$ if we can get a good estimate on $P(\mathbf{c}_i|\mathbf{x})$

k-Nearest Neighbor


- How would we estimate $P(\mathbf{c}_i / \mathbf{x})$ from a set of n labeled samples?

- Recall our estimate for density: $p(\mathbf{x}) \approx \frac{k/n}{V}$

- Let's place a cell of volume V around \mathbf{x} and capture k samples

- k_i samples amongst k labeled \mathbf{c}_i then:

$$p(\mathbf{c}_i, \mathbf{x}) \approx \frac{k_i / n}{V}$$



- Using conditional probability, let's estimate posterior:

$$p(\mathbf{c}_i | \mathbf{x}) = \frac{p(\mathbf{x}, \mathbf{c}_i)}{p(\mathbf{x})} = \frac{p(\mathbf{x}, \mathbf{c}_i)}{\sum_{j=1}^m p(\mathbf{x}, \mathbf{c}_j)} \approx \frac{k_i / n}{V \sum_{j=1}^m \frac{k_j / n}{V}} = \frac{k_i}{\sum_{j=1}^m k_j} = \frac{k_i}{k}$$

k-Nearest Neighbor


- Thus our estimate of posterior is just the fraction of samples which belong to class \mathbf{c}_i :

$$p(\mathbf{c}_i | \mathbf{x}) = \frac{k_i}{k}$$

- This is a very simple and intuitive estimate
- Under the zero-one loss function (MAP classifier) just choose the class which has the largest number of samples in the cell
- Interpretation is: given an unlabeled example (that is \mathbf{x}), find k most similar labeled examples (closest neighbors among sample points) and assign the most frequent class among those neighbors to \mathbf{x}

k-Nearest Neighbor: Example

- Back to fish sorting
 - Suppose we have 2 features, and collected sample points as in the picture
 - Let $k = 3$



- 2 sea bass, 1 salmon are the 3 nearest neighbors
- Thus classify as sea bass

kNN: How Well Does it Work?

- kNN rule is certainly simple and intuitive, but does it work?
- Pretend that we can get an unlimited number of samples
- By definition, the best possible error rate is the Bayes rate E^*
- Even for $k=1$, the nearest-neighbor rule leads to an error rate greater than E^*
- But as $n \rightarrow \infty$, it can be shown that nearest neighbor rule error rate is smaller than $2E^*$
- If we have a lot of samples, the kNN rule will do very well !

1NN: Voronoi Cells





FIGURE 4.13. In two dimensions, the nearest-neighbor algorithm leads to a partitioning of the input space into Voronoi cells, each labeled by the category of the training point it contains. In three dimensions, the cells are three-dimensional, and the decision boundary resembles the surface of a crystal. From: Richard O. Duda, Peter E. Hart, and David G. Stork, *Pattern Classification*. Copyright © 2001 by John Wiley & Sons, Inc.

*k*NN: Multi-Modal Distributions

- Most parametric distributions would not work for this 2 class classification problem:
- Nearest neighbors will do reasonably well, provided we have a lot of samples




***k**NN: How to Choose **k**?*

- In theory, when the infinite number of samples is available, the larger the **k**, the better is classification (error rate gets closer to the optimal Bayes error rate)
- But the caveat is that all **k** neighbors have to be close to **x**
 - Possible when infinite # samples available
 - Impossible in practice since # samples is finite

***k**NN: How to Choose **k**?*

- In practice
 - 1.** **k** should be large so that error rate is minimized
 - **k** too small will lead to noisy decision boundaries
 - 2.** **k** should be small enough so that only nearby samples are included
 - **k** too large will lead to over-smoothed boundaries
- Balancing **1** and **2** is not trivial
 - This is a recurrent issue, need to smooth data, but not too much

*k*NN: How to Choose *k*?




- For $k = 1, \dots, 7$ point \mathbf{x} gets classified correctly
 - red class
- For larger k classification of \mathbf{x} is wrong
 - blue class

kNN: Computational Complexity

- Basic ***kNN*** algorithm stores all examples. Suppose we have ***n*** examples each of dimension ***k***
 - $O(d)$ to compute distance to one example
 - $O(nd)$ to find one nearest neighbor
 - $O(knd)$ to find ***k*** closest examples
 - Thus complexity is $O(knd)$
- This is prohibitively expensive for large number of samples
- But we need large number of samples for ***kNN*** to work well!

Reducing Complexity: Editing 1NN


- If all voronoi neighbors have the same class, a sample is useless, we can remove it:



- Number of samples decreases
- We are guaranteed that the decision boundaries stay the same

Reducing Complexity: *k*NN prototypes

- Explore similarities between samples to represent data as search trees of *prototypes*



- Advantages: Complexity decreases
- Disadvantages:
 - finding good search tree is not trivial
 - will not necessarily find the closest neighbor, and thus **not** guaranteed that the decision boundaries stay the same


kNN: Selection of Distance

- So far we assumed we use Euclidian Distance to find the nearest neighbor:

$$D(\mathbf{a}, \mathbf{b}) = \sqrt{\sum_k (\mathbf{a}_k - \mathbf{b}_k)^2}$$

- However some features (dimensions) may be much more discriminative than other features (dimensions)
- Euclidean distance treats each feature as equally important

*k*NN: Extreme Example of Distance Selection



- decision boundaries for blue and green classes are in red
- These boundaries are really bad because
 - feature 1 is discriminative, but its scale is small
 - feature 2 gives no class information (noise) but its scale is large

*k*NN: Selection of Distance

- Extreme Example
 - feature 1 gives the correct class: 1 or 2
 - feature 2 gives irrelevant number from 100 to 200
- Suppose we have to find the class of $x=[1 \ 100]$ and we have 2 samples $[1 \ 150]$ and $[2 \ 110]$


$$D\left(\begin{bmatrix} 1 \\ 100 \end{bmatrix}, \begin{bmatrix} 1 \\ 150 \end{bmatrix}\right) = \sqrt{(1-1)^2 + (100-150)^2} = 50 \quad D\left(\begin{bmatrix} 1 \\ 100 \end{bmatrix}, \begin{bmatrix} 2 \\ 110 \end{bmatrix}\right) = \sqrt{(1-2)^2 + (100-110)^2} = 10.5$$

- $x = [1 \ 100]$ is misclassified!
- The denser the samples, the less of the problem
 - But we rarely have samples dense enough

kNN: Selection of Distance

- Notice the 2 features are on different scales:
 - feature 1 takes values between 1 or 2
 - feature 2 takes values between 100 to 200
- We could normalize each feature to be between of mean 0 and variance 1
- If \mathbf{X} is a random variable of mean μ and variance σ^2 , then $(\mathbf{X} - \mu)/\sigma$ has mean 0 and variance 1
- Thus for each feature vector \mathbf{x}_i , compute its sample mean and variance, and let the new feature be $[\mathbf{x}_i - \text{mean}(\mathbf{x}_i)]/\text{sqrt}[\text{var}(\mathbf{x}_i)]$
- Let's do it in the previous example

*k*NN: Normalized Features



- The decision boundary (in red) is very good now!

***k*NN: Selection of Distance**

- However in high dimensions if there are a lot of irrelevant features, normalization will not help

$$D(\mathbf{a}, \mathbf{b}) = \sqrt{\sum_k (\mathbf{a}_k - \mathbf{b}_k)^2} = \sqrt{\sum_i (\mathbf{a}_i - \mathbf{b}_i)^2 + \sum_j (\mathbf{a}_j - \mathbf{b}_j)^2}$$

discriminative *noisy*
feature *features*

- If the number of discriminative features is smaller than the number of noisy features, Euclidean distance is dominated by noise

kNN: Feature Weighting

- Scale each feature by its importance for classification

$$D(\mathbf{a}, \mathbf{b}) = \sqrt{\sum_k \mathbf{w}_k (\mathbf{a}_k - \mathbf{b}_k)^2}$$

- Can learn the weights \mathbf{w}_k from the training data
 - Increase/decrease weights until classification improves

kNN Summary

- Advantages
 - Can be applied to the data from any distribution
 - Very simple and intuitive
 - Good classification if the number of samples is large enough
- Disadvantages
 - Choosing best k may be difficult
 - Computationally heavy, but improvements possible
 - Need large number of samples for accuracy
 - Can never fix this without assuming parametric distribution