Article Submitted to Journal of Symbolic Computation

Computation of Canonical Forms for Ternary Cubics

Irina A. Kogan[1] and Marc Moreno Maza[2,3]

[1] *Yale University, Dept of Math, New Haven, CT 06520-8283, USA*
irina.kogan@yale.edu

[2] *Université de Lille 1, LIFL, 59655 Villeneuve d'Ascq France*
moreno@lifl.fr

[3] *University of Western Ontario, ORCCA, London, ON N6A 5B7, Canada*
moreno@orcca.on.ca

Abstract

In this paper we conduct a careful study of the equivalence classes of ternary cubics under general linear changes of variables. Our new results are based on the method of moving frames and involve triangular decompositions of algebraic varieties. We provide a computationally efficient algorithm that matches an arbitrary ternary cubic with its canonical form and explicitly computes a corresponding linear change of coordinates. We separately consider complex and real transformations. We also describe a classification of the symmetry groups of ternary cubics.

Introduction.

One of the central problems of classical invariant theory is the equivalence and symmetry of multivariable polynomials under linear changes of variables. The standard action of the general linear group on the m dimensional complex linear space $\mathbb{C}^m$ induces an action $F \to \bar{F}$ on the linear space of polynomials $\mathbb{C}[\mathbf{x}]$ defined by

$$\bar{F}(g \cdot \mathbf{x}) = F(\mathbf{x}), \tag{1}$$

where $g \in GL(m, \mathbb{C})$, $\mathbf{x} \in \mathbb{C}^m$ and $F, \bar{F} \in \mathbb{C}[\mathbf{x}]$ are polynomials in m variables.

The set of homogeneous polynomials of degree n is mapped to itself under the $GL(m, \mathbb{C})$ action defined by (1). Two polynomials are said to be equivalent if there is a linear change of variables that transforms one into the other. The same problem can be considered over the field of real numbers. The group of symmetries of a polynomial consists of all linear changes of variables that leave the polynomial unchanged. It is desirable to describe the classes of equivalent

polynomials, or, in other words, classify the orbits of the action of a general linear group on the linear space of polynomials.

In this paper we consider *ternary cubics*, that is, homogeneous polynomials in three variables of degree three. Their classification is well known, see for instance [Gurevich, 1964], [Kraft, 1985], [Bix, 1998]. In each orbit we identify a "simple" canonical form and provide an algorithm, that matches an arbitrary cubic with its canonical form. A corresponding linear change of variables is computed explicitly. Note that the most naive approach to the latter problem, solving (1) for the group parameters, requires very hard computations. Our algorithm is based on the novel differential geometry approach first introduced by Olver [1999]. The solution of the equivalence problem over real linear transformations can be approached by similar techniques but is more subtle. We indicate how to adapt our new methods to the real case in Section 6.

Polynomials of degree n in m variables form a linear space W of dimension $\binom{m+n-1}{n}$ isomorphic to the n-th symmetric tensor power of $\mathbb{C}^m$. The coefficients of polynomials can serve as coordinates on this space and formula (1) induces an irreducible representation of $GL(m,\mathbb{C})$ on W. One can try to classify polynomials by computing continuous *invariants*, that is, functions $H(\dots, a_{i_1,\dots,i_m}, \dots)$ in the coefficients of a polynomial, invariant under the action (1). However, the action (1) is not regular: the dimensions of the orbits vary and all orbits, except the orbit of the zero polynomial, are not closed. Hence the orbits can not be distinguished by continuous invariants. More information can be obtained from *covariants*, $H(\mathbf{x}, \dots, a_{i_1,\dots,i_m}, \dots)$, continuous functions that are invariant under the simultaneous action of the general linear group on the variables and the coefficients of polynomials. The simplest example of a covariant is the polynomial $F(\mathbf{x})$ itself (see formula (1)). Computation of *rational* invariants and covariants, as well as polynomial relations, or *sysygies*, among them is the main subject of the nineteenth century classical theory. The overview of the classical methods for constructing rational covariants as well their application to the classification of polynomials can be found in [Grace and Young, 1903], [Gurevich, 1964], [Olver, 1999]. A set of rational covariants is called *fundamental* if any other rational covariant can be expressed as a rational function of the fundamental ones. The existence of a finite fundamental set of covariants follows from the finite basis theorem for the actions of linear reductive groups proved by Hilbert in 1890 [Ackerman and R., 1978]. The number of fundamental covariants, however, grows dramatically, with the increase of the degree of polynomials even for a fixed number of variables. The Hilbert's proof of the finite basis theorem became a turning point from classical computational approach in the invariant theory to its modern formulation in terms of algebraic geometry [Mumford, 1994], [Kraft, 1985], [Vinberg and Popov, 1989] and representation theory [Howe, 1989].

Despite the enormous amount of work devoted to the subject, the general classification of homogeneous polynomials and their symmetry groups remains unknown even for the case of polynomials in two or three variables (over $\mathbb{C}$ or $\mathbb{R}$) except for polynomials of low degree.

The main idea of moving frame approach, appeared in [Olver, 1999], is to consider the graph of a polynomial $F(\mathbf{x})$ in m variables as a submanifold in $(m+1)$-dimensional complex (or real) space therefore reducing the question to equivalence problems for submanifolds. The latter problem can be resolved by Cartan's method of moving frames, which involves computation of a fundamental set of differential invariants. We note that, in contrast with the classical approach, a set of differential invariants is called *fundamental* if any other differential invariant can be at least locally expressed as a smooth function of the fundamental ones. Fundamental differential invariants parameterize the *signature manifold*, or the *classifying manifold*, of F [Olver, 1995]. Two equivalent polynomials have the same signature manifold. Moreover the signatures can be used to determine the symmetry groups (see Section 4) and to describe the geometry of the orbits for the $GL(3,\mathbb{C})$-action on the linear space of polynomials (see Section 3).

In the joint paper by Peter Olver and the first author [Berchenko and Olver, 2000] the moving frame method has been applied to polynomials in two variables (binary forms). A MAPLE package which determines the dimension of the symmetry group of a given polynomial and, in the case when the symmetry group is finite, computes it explicitly has been provided. Computing differential invariants becomes more challenging for the homogeneous polynomials in more than two variables, even in the next case of polynomials in three variables. This work has been completed by the first author in her thesis [Kogan, 2000], using the inductive moving frame construction presented in [Kogan, 2001]. These invariants were used to classify the symmetry groups for ternary cubics and to solve some equivalence problems. For instance, a necessary and sufficient condition for a homogeneous polynomial in three variables to be equivalent to $x^n + y^n + z^n$ was computed in [Kogan, 2000] and [Schost, 2001]. The general problem of expressing a form as a sum of powers of linear forms has applications in signal processing and is discussed in [Common and Mourrain, 1996], [Reznick, 1992].

To the advantage of the moving frames approach, it is applicable to polynomials of any degree in any number of variables, as well as one can consider various group actions. Moreover, one can use the same invariants for polynomials of any degree, as long as the number of variables and the group action remain the same. In practice, however, we are confronted with challenging elimination problems for multivariable polynomials, not solvable by the standard algorithm based on Gröbner bases computations. The results for the case of ternary cubics, presented here, were obtained by using triangular decompositions. More precisely, the signature manifolds of the canonical ternary forms were obtained by means of the PARDI algorithm by Boulier, Lemaire and the second author [Boulier et al., 2001a]. The justification of these computations follows from the theory of regular chains introduced by Kalkbrener [1993] and its recent developments [Aubry et al., 1999], [Boulier et al., 2001b]. The computations were performed by the ALDOR [Bronstein et al., 2001] implementation of the PARDI algorithm written by the second author.

Comparing to the preliminary version of this paper [Kogan and Moreno Maza, 2002] we present here two recent developments of our work. First we discuss in deep the computation of a linear transformation from an arbitrary ternary cubic into its canonical form. We show that this can be obtained with very simple calculations (resultant/gcd of bivariate polynomials with small total degree and factorization of univariate polynomials of small degree). Secondly, we present a classification of real ternary cubics under linear transformation over real numbers and discuss how to apply the moving frame method in order to match a real cubic with its real class. We note that, as in the complex case, the real classification of the cubics is known. Our goal was to confirm the efficiency of the moving frame method, which can be applied to polynomials of higher degree.

The paper is organized as follows.

Section 1 contains a formulation of the equivalence and symmetry problems for multivariable polynomials under linear changes of variables together with a description of the equivalence classes of ternary cubics under this action.

Section 2 contains a derivation of a moving frame and the corresponding differential invariants for the action considered in the previous section. These invariants are not rational. However we use them to build a new set of fundamental rational invariants which leads to exceptionally simple signature manifolds.

Section 3 describes the signature manifolds for each of the equivalence classes. We also make some observations about the geometry of the orbits.

Section 4 contains an algorithm for computing the symmetry groups listed in the previous section.

Section 5 provides an algorithm that matches an arbitrary ternary cubic with its canonical form and computes a corresponding linear change of coordinates.

Section 6 discusses a classification of real ternary cubics under linear transformation over real numbers.

1. Equivalence and Symmetry.

Let us consider the action of the general linear group on the space of polynomials over $\mathbb{C}$ in m variables:

$$\bar{F}(g \cdot \mathbf{x}) = F(\mathbf{x}), \text{ equivalently } \bar{F}(\mathbf{x}) = F(g^{-1}\mathbf{x}) \tag{2}$$

where $g \in GL(m, \mathbb{C})$, $\mathbf{x} \in \mathbb{C}^m$ and $F, \bar{F} \in \mathbb{C}[\mathbf{x}]$.

Definition 1.1: *A polynomial F is said to be equivalent to a polynomial $\bar{F}$ if there exists $g \in GL(m, \mathbb{C})$ such that $\bar{F}(g \cdot \mathbf{x}) = F(\mathbf{x})$. In this case we will also write $\bar{F} = g \cdot F$*

EXAMPLE 1.1: *The polynomial* $x^3+y^3+z^3$ *is equivalent to* $x^3+6\,x\,y^2-6\,x\,y\,z-3\,y^2\,z+3\,y\,z^2$ *under the transformation*

$$x \mapsto x-y; \quad y \mapsto x+y-z; \quad z \mapsto z-x.$$

DEFINITION 1.2: *The symmetry group of* F *is the subgroup* $G_F \subset GL(m,\mathbb{C})$ *consisting of all linear transformations that map* F *to itself.*

EXAMPLE 1.2: *The connected component of the symmetry group of the cubic* $z\,(x^2+y\,z)$ *is a two-dimensional noncommutative (affine) group, generated by two one parametric subgroups:*

$$\begin{pmatrix} 1 & 0 & \alpha \\ -2\alpha & 1 & -\alpha^2 \\ 0 & 0 & 1 \end{pmatrix} \quad \textit{and} \quad \begin{pmatrix} \beta & & \\ & \beta^4 & \\ & & \beta^{-2} \end{pmatrix}.$$

We note that the symmetry groups of equivalent polynomials are related by matrix conjugation:

$$\bar{F}(\mathbf{x}) = F(g\cdot\mathbf{x}) \Longrightarrow G_{\bar{F}} = g G_F g^{-1}. \tag{3}$$

Thus the problem of the classification of polynomials is closely related to the problem of the classification of their symmetry groups up to matrix conjugation. In Section 4 we describe an algorithm (first appeared in [Olver, 1999] and used in [Kogan, 2000]) to determine the dimension of the symmetry group of a given polynomials and, in the case when the cardinality of the symmetry group is finite, to compute its order.

We focus on homogeneous polynomials of degree three with three variables, also called ternary cubics. We reproduce below the known classification of ternary cubics up to a linear transformation. We start with irreducible cubics. A nonsingular projective variety V defined by an irreducible ternary cubic $F(x,y,z)$ over $\mathbb{C}$ is an *elliptic curve*. The study of elliptic curves over complex, real and rational numbers is a very interesting and important problem in number theory with applications to the theory of modular forms [Knapp, 1992], [Lang, 1987]. Their canonical forms, also called Weierstrass normal forms, can be obtained by transforming one of the inflection points of V to the infinite point $(0,1,0)$, and the tangent line at this point to the line $(k,1,0)$ at infinity. A complex cubic has 9 inflection points, that can be found by computing the common roots of the cubic itself and its Hessian [Knapp, 1992], [Bix, 1998].

THEOREM 1.1: *An irreducible cubic* $F(x,y,z)$ *can be transformed under a complex linear change of variables to one of the following forms:*

1 *If* $F(x,y,z)$ *defines a nonsingular projective variety then it is either equivalent to:*

a) *a cubic in the one-parametric family:*

$$x^3+\mu\,x\,z^2+z^3-y^2\,z,$$

where $\mu \neq 0$ (otherwise it is an irreducible cubic of type c) below) and $\mu^3 \neq -27/4$ (otherwise it is equivalent to the reducible cubic of type b) below).
or either to

$$\textbf{b)} \quad x^3 + x\,z^2 - y^2\,z \quad \textit{or} \quad \textbf{c)} \quad x^3 + z^3 - y^2\,z.$$

2. *If $F(x, y, z)$ defines a singular projective variety then it is equivalent to either*

$$\textbf{a)} \quad x^3 - y^2\,z \quad \textit{or} \quad \textbf{b)} \quad x^2(x+z) - y^2\,z.$$

THEOREM 1.2: *A reducible cubic $F(x, y, z)$ is equivalent under a linear change of variables to one of the following forms:*

1. *If it is a product of quadratic and linear factors then it is equivalent to either*

$$\textbf{a)} \quad z\,(x^2 + yz) \qquad \textit{or} \qquad \textbf{b)} \quad z\,(x^2 + y^2 + z^2).$$

2. *If it is a product of three linear factors and*

- **a)** *three factors are linearly independent factors, then the cubic is equivalent to xyz.*
- **b)** *three factors are linearly dependent, but any pair of them is linearly independent, then the cubic is equivalent to $xy(x + y)$;*
- **c)** *two factors are the same, then the cubic is equivalent to x^2y;*
- **d)** *all three factors are the same, then it is equivalent to x^3.*

The classification of reducible cubics can be obtained by very elementary methods [Kogan, 2000].

In the next two sections, we address the problem of matching a cubic F with one of the canonical forms listed above. Once the canonical form $\bar{F}$ is identified, we would like to determine a linear transformation such that $\bar{F} = F(g \cdot \mathbf{x})$. The most straightforward approach, solving the equation

$$\bar{F}(x, y, z) = F\left(\alpha x + \beta y + \lambda z\,, \gamma x + \delta y + \mu z\,, ax + by + \eta z\right).$$

for the group parameters, turns out to be computationally impractical. A classical alternative to our algorithm is the constructive approach described [Bix, 1998], that leads to the above classification. One of the advantages of our algorithm is its generality, it is applicable to polynomials of higher degree and larger number of variables. Another advantage, is that one can instantaneously determine the canonical form of a given polynomial, without computing a corresponding transformation. In Section 5 we propose a practical algorithm to find a transformation that bring a cubic to its canonical form. We show that its efficiency compares to the classical algorithm described in [Bix, 1998].

2. Differential Invariants and Signature Manifolds.

The classification algorithms presented in the next sections are based on the moving frame method first developed by Cartan [1935] for the solution of equivalence and symmetry problems in differential geometry. We refer the reader to the above reference as well as to [Guggenheimer, 1963] for the classical formulation of the method, and to [Fels and Olver, 1999], [Griffiths, 1974], [Green, 1978] for its modern algorithmic formulation. In this section we implement the inductive variation [Kogan, 2001] of the moving frame algorithm, presented in [Fels and Olver, 1999], to compute a fundamental set of differential invariants for ternary cubics.

Following the ideas of [Olver, 1999], we consider the graph of a homogeneous cubic polynomial $u = F(x, y, z)$ as a smooth three-dimensional submanifold in $\mathbb{C}^4$. Coordinate functions in $\mathbb{C}^4$ are chosen to be (x, y, z, u). A group element

$$g = \begin{pmatrix} \alpha & \beta & \lambda \\ \gamma & \delta & \mu \\ a & b & \eta \end{pmatrix} \in GL(3, \mathbb{C}) \tag{4}$$

acts on $\mathbb{C}^4$ by leaving coordinate u unchanged and transforming linearly x, y, z:

$$\begin{array}{ccc} u & \to & u\,, \\ \begin{pmatrix} x \\ y \\ z \end{pmatrix} & \to & \begin{pmatrix} \alpha & \beta & \lambda \\ \gamma & \delta & \mu \\ a & b & \eta \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}. \end{array} \tag{5}$$

The graph Γ_F of $u = F(x, y, z)$ is transformed under this action to the graph $g \cdot \Gamma_F$ of

$$u = F\left(\tilde{\alpha}\, x + \tilde{\beta}\, y + \tilde{\lambda}\, z\,, \tilde{\gamma}\, x + \tilde{\delta}\, y + \tilde{\mu}\, z\,, \tilde{a}\, x + \tilde{b}\, y + \tilde{\eta}\, z\right), \tag{6}$$

where $\tilde{\alpha}, \ldots, \tilde{\eta}$ denote the entries of the matrix g^{-1}. In order to reduce the dimension of the jet spaces which will be introduced below, we prefer to work with inhomogeneous (projective) version of F, the unique polynomial in two variables p, q defined by:

$$f(p, q) = F(p, q, 1). \tag{7}$$

For example, the polynomial $f(p, q) = pq$ is the inhomogeneous version of the cubic $F(x, y, z) = xyz$. We note that the degree of f is less than or equal to three, but we keep referring to f as a cubic, since the knowledge of the initial degree is necessary to restore F from f:

$$F(x, y, z) = z^3\, f\left(\frac{x}{z}, \frac{y}{z}\right).$$

We consider the graph Γ_f of the polynomial $u = f(p, q)$ as a surface in the three

dimensional complex space. From (6) and (7) we deduce that Γ_f is transformed to the graph of:

$$u = \left(\tilde{a}\,p + \tilde{b}\,q + \tilde{\eta}\right)^3 \; f\left(\frac{\tilde{\alpha}\,p + \tilde{\beta}\,q + \tilde{\lambda}}{\tilde{a}\,p + \tilde{b}\,q + \tilde{\eta}}, \frac{\tilde{\gamma}\,p + \tilde{\delta}\,q + \tilde{\mu}}{\tilde{a}\,p + \tilde{b}\,q + \tilde{\eta}}\right). \tag{8}$$

This corresponds to the following local action of $GL(3,\mathbb{C})$ on $\mathbb{C}^3$, defined at each point for the group elements that do not allow the denominator vanish.

$$\begin{aligned} u &\rightarrow \bar{u} = (a\,p + b\,q + \eta)^{-3}\; u, \\ p &\rightarrow \bar{p} = \frac{\alpha\,p + \beta\,q + \lambda}{a\,p + b\,q + \eta}, \\ q &\rightarrow \bar{q} = \frac{\gamma\,p + \delta\,q + \mu}{a\,p + b\,q + \eta}. \end{aligned} \tag{9}$$

The application of the moving frame method is justified by the following simple proposition:

PROPOSITION 2.1: *Two homogeneous cubics $F(x,y,z)$ and $\bar{F}(x,y,z)$ are equivalent under a linear change of variables (see Definition 1.1) if and only if the graph $u = F(x,y,z)$ in $\mathbb{C}^4$ can be mapped to the graph $u = \bar{F}(x,y,z)$ under the transformation (5) or, equivalently, the graph $u = f(p,q)$ in $\mathbb{C}^3$ of the inhomogeneous version of F can be mapped to $u = \bar{f}(p,q)$ by (9).*

The first step is to prolong the graph $u = f(p,q)$ and the group action to the jet space of smooth functions in two variables. In our case the n-th jet space J^n is a (n^2+3n+6)-dimensional complex space parameterized by coordinates $(p, q, u, u_{1,0}, u_{0,1}, \ldots, u_{i,j}, \ldots u_{0,n})$, where $i+j \leq n$ and $u_{i,j}$ are formal coordinates that correspond to all possible partial derivatives of u with respect to p and q. Given a particular polynomial $f(p,q)$, one can actually compute these derivatives in order to obtain a two-dimensional surface in J^n:

$$u = f(p,q), \quad u_{i,j} = \frac{\partial^{i+j} u}{\partial^i p \partial^j q}, \tag{10}$$

which is called n-th prolongation of the graph $u = f(p,q)$ and is denoted by $\Gamma_f^{(3)}$. We note that since f is a polynomial of degree 3, all the derivatives of order higher than three are zero, and thus one can solve the equivalence problem on the third order jet space J^3 of dimension 12.

REMARK 2.1: *Not every two-dimensional surface in J^3 corresponds to a prolongation of a cubic polynomial f. In fact there is one and only one such graph passing through every point of J^3. Remarkably, when polynomials of an arbitrary degree n are considered, one does not need to prolong the graph up to the order n to solve the equivalence and symmetry problems, but at most up to the order 6. Although there are several prolonged graphs of polynomials of degree > 6 passing*

through a point in J^6, there is at most one polynomial from each equivalence class passing through each point. That is, if $\Gamma_f^{(6)}$ and $\Gamma_{g\cdot f}^{(6)}$ are passing through the same point in J^6, then $f = g \cdot f$ (see rigidity theorems [Fels and Olver, 1999] Sect. 14).

The next step is to prolong the action (9) to the jet space J^3, that is, to define a transformation for the coordinates $u_{i,j}$, in a way that it commutes with the process of the prolongation of the graph (10). According to [Fels and Olver, 1999] and Chapter 2 of [Olver, 1986], this can be done in the unique way:

$$u_{i,j} \to D_p^i \, D_q^j(\bar{u}),$$

where D_p and D_q are differential operators defined by:

$$D_p = C\left((p\,A + \delta\,\eta - b\,\mu)\,\frac{d}{dp} + (q\,A + a\,\mu - \gamma\eta)\,\frac{d}{dq}\right),$$
$$D_q = C\left((p\,B + \beta\,\eta - b\,\lambda)\,\frac{d}{dp} + (q\,B + a\,\lambda - \alpha\eta)\,\frac{d}{dq}\right),$$

where $C = \frac{a\,p+b\,q+\eta}{\det(g)}$, $A = \delta\,a - \gamma\,b$ and $B = \beta\,a - \alpha\,b$.

For example, $u_{1,0}$ is transformed to

$$D_p(\bar{u}) = \frac{-3A\,u + (p\,A + \delta\,\eta - b\,\mu)\;u_{1,0} + (q\,A + a\,\mu - \gamma\eta)\;u_{0,1}}{(a\,p + b\,q + \eta)^2\,\det(g)}.$$

To avoid long and, in fact, unnecessary formulae we will denote the rest of the transformations symbolically:

$$u_{i,j} \to \bar{u}_{i,j} = \Phi_{i,j}(\mathbf{u}^{(3)}, g), \tag{11}$$

where $0 \le i + j \le 3$ and $\Phi_{i,j}$ are smooth functions of the group parameters (4) and coordinates of J^3.

An important observation is that the action of the group $G = GL(3,\mathbb{C})$ is *locally free* on J^3, which means that the stabilizers of each points are discrete. In fact, the isotropy group of a generic point consists of three matrices in the complex case and of identity matrix only in the real case. It follows that each orbit is locally diffeomorphic to G and thus has dimension equal to $\dim G = 9$. One can find a *cross-section* $\mathcal{K}$ to the orbits, that is, a submanifold in J^3 of dimension $3 = \dim J^3 - \dim G$, which is transversal to the orbits and intersects each orbit in an open subset of J^3 once. It turns out that the submanifold

$$\mathcal{K} = \{\mathbf{u}^{(3)} \in J^3 | p = 0, q = 0, u = 1, \tag{12}$$
$$u_{1,0} = 0, u_{0,1} = 0, u_{2,0} = 0, u_{0,2} = 0, u_{3,0} = 1, u_{0,3} = 1\}$$

can serve this purpose on a dense subset of J^3. Computationally it is reflected in the existence of a finite set of the solutions of the following system (see (9,11)):

$$\bar{p} = 0,\ \bar{q} = 0,\ \bar{u} = 1,\ \Phi_{1,0} = 0,\ \Phi_{0,1} = 0, \tag{13}$$
$$\Phi_{2,0} = 0,\ \Phi_{0,2} = 0,\ \Phi_{3,0} = 1,\ \Phi_{0,3} = 1.$$

for the group parameters at almost every point of J^3. A solution for this system yields a matrix $g(\mathbf{u}^{(3)})$ which maps a point $\mathbf{u}^{(3)} \in J^3$ to $\mathcal{K}$ under the prolonged transformation (9, 11). The matrix-valued function $g(\mathbf{u}^{(3)})$, defined at almost each point of J^3, is called a *moving frame* [Fels and Olver, 1999]. By construction the map $\pi(\mathbf{u}^{(3)}) = g(\mathbf{u}^{(3)}) \cdot \mathbf{u}^{(3)}$ defines a projection π of J^3 onto $\mathcal{K}$ along the orbits. In particular, one can restrict π to the prolonged graph of $u = f(p, q)$ defined by (10), to obtain a projection $\pi_f : \Gamma_f \to \mathcal{K}$. As the following two theorems indicate the image of Γ_f in $\mathcal{K}$, called the *signature manifold* of f, provides the key to the solution of the symmetry and equivalence problems.

DEFINITION 2.1: *The signature manifold $\mathcal{C}_f$ of a cubic polynomial $f(p, q)$ is the projection along the orbits of the third prolongation (10) of its graph $\Gamma_f^{(3)}$ to the cross-section $\mathcal{K}$.*

THEOREM 2.1: *Two cubics $f(p, q)$ and $\bar{f}(p, q)$ are equivalent under a linear change of variables if and only if their signature manifolds coincide, $\mathcal{C}_f = \mathcal{C}_{\bar{f}}$.*

Figure (1) shows a graphical sketch of the moving frame method. For the ab-



Figure 1: Construction of the signature manifold.

solute majority of cubic polynomials the projections of their two-dimensional graphs to $\mathcal{K}$ remains two-dimensional, but some of the cubics, whose graphs are

not transversal to the orbits, will have the projections (signatures) of a smaller dimension. The drop in the dimension of the signature manifold reflects the increase in the dimension of the symmetry group of f.

THEOREM 2.2: *The dimension of the symmetry group G_f of $f(p,q)$ equals to* $\dim \Gamma_f - \dim \mathcal{C}_f = 2 - \dim \mathcal{C}_f$.

See [Fels and Olver, 1999] (Theorems 14.7 and 14.9) and [Olver, 1995] (Theorems 8.19 and 8.22) for the proofs of Theorems 2.1 and 2.2 above. The computation of the signatures relies on the solution of the system (13) for the group parameters. However, the formulae for $\Phi_{i,j}$ in the second or third order are too long and frightening to approach. We can, however, split the problem into three feasible sub-problems, by decomposing the matrix of g into a product of three matrices $g = LRT$, where

$$L = \begin{pmatrix} \alpha & \beta & 0 \\ \gamma & \delta & 0 \\ 0 & 0 & 1 \end{pmatrix}, R = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ a & b & c \end{pmatrix}, T = \begin{pmatrix} 1 & 0 & t \\ 0 & 1 & s \\ 0 & 0 & 1 \end{pmatrix}.$$

We apply the inductive construction of moving frames [Kogan, 2001], [Kogan, 2000] to this product. The resulting differential invariants are valid for ternary forms of *any* degree n, with $n = 3$ in the case of cubics.

The product of matrices RT with

$$t = -p,\ s = -q,\ c = u^{1/n},\ a = \frac{u_{1,0}}{n\, u^{\frac{n-1}{n}}},\ b = \frac{u_{0,1}}{n\, u^{\frac{n-1}{n}}} \tag{14}$$

transforms a point $\mathbf{u}^{(3)}$ to a point on the cross-section $\mathcal{K}_1 = \{\mathbf{u}^{(3)} \in J^3 | p = 0, q = 0, u = 1, u_{1,0} = 0, u_{0,1} = 0\} \supset \mathcal{K}$. Coordinates, $u_{2,0}, u_{0,2}, u_{3,0}, u_{2,1}, u_{1,2}, u_{0,3}$ are transformed to the following expressions:

$$\begin{aligned}
Q_{2,0} &= u^{-\frac{2(n-1)}{n}} \left[u_{2,0}\, u - \frac{(n-1)}{n} u_{1,0}^2 \right] = u^{-\frac{2(n-1)}{n}} \hat{Q}_{2,0}, \\
Q_{1,1} &= u^{-\frac{2(n-1)}{n}} \left[u_{1,1}\, u - \frac{(n-1)}{n} u_{1,0} u_{0,1} \right] = u^{-\frac{2(n-1)}{n}} \hat{Q}_{1,1}, \\
Q_{0,2} &= u^{-\frac{2(n-1)}{n}} \left[u_{0,2}\, u - \frac{(n-1)}{n} u_{0,1}^2 \right] = u^{-\frac{2(n-1)}{n}} \hat{Q}_{0,2}, \\
Q_{3,0} &= u^{-\frac{3(n-1)}{n}} \left[u_{3,0}\, u^2 - 3\frac{(n-2)}{n} u_{2,0} u_{1,0}\, u + 2\frac{(n-1)(n-2)}{n^2} u_{1,0}^3 \right], \\
Q_{2,1} &= u^{-\frac{3(n-1)}{n}} \left[u_{2,1}\, u^2 - \frac{(n-2)}{n} (u_{2,0} u_{0,1} + 2\, u_{1,1} u_{1,0})\, u + 2\frac{(n-1)(n-2)}{n^2} u_{1,0}^2 u_{0,1} \right], \\
Q_{1,2} &= u^{-\frac{3(n-1)}{n}} \left[u_{1,2}\, u^2 - \frac{(n-2)}{n} (u_{0,2} u_{1,0} + 2\, u_{1,1} u_{0,1})\, u + 2\frac{(n-1)(n-2)}{n^2} u_{0,1}^2 u_{1,0} \right], \\
Q_{0,3} &= u^{-\frac{3(n-1)}{n}} \left[u_{0,3}\, u^2 - 3\frac{(n-2)}{n} u_{0,2} u_{0,1}\, u + 2\frac{(n-1)(n-2)}{n^2} u_{0,1}^3 \right],
\end{aligned} \tag{15}$$

Symbols $\hat{Q}_{2,0}, \hat{Q}_{1,1}, \hat{Q}_{0,2}$ are introduced to be used later. By restricting the action of L to $\mathcal{K}_1$ we obtain the following transformations of $Q_{i,j}$:

$$\begin{aligned}
\bar{Q}_{2,0} &= \frac{1}{A^2}\left(\delta^2 Q_{2,0} - 2\gamma\delta Q_{1,1} + \gamma^2 Q_{0,2}\right), \\
\bar{Q}_{1,1} &= \frac{1}{A^2}\left(-\delta\beta Q_{2,0} + (\gamma\beta + \alpha\delta) Q_{1,1} - \alpha\gamma Q_{0,2}\right), \\
\bar{Q}_{0,2} &= \frac{1}{A^2}\left(\beta^2 Q_{2,0} - 2\alpha\beta Q_{1,1} + \alpha^2 Q_{0,2}\right), \\
\bar{Q}_{3,0} &= \frac{1}{A^3}\left(\delta^3 Q_{3,0} - 3\gamma\delta^2 Q_{2,1} + 3\gamma^2\delta Q_{1,2} - \gamma^3 Q_{0,3}\right), \\
\bar{Q}_{2,1} &= \frac{1}{A^3}\left(-\delta^2\beta Q_{3,0} + \delta(2\gamma\beta + \alpha\delta) Q_{2,1} - \gamma(\gamma\beta + 2\alpha\delta) Q_{1,2} + \alpha\gamma^2 Q_{0,3}\right) \\
\bar{Q}_{1,2} &= \frac{1}{A^3}\left(\delta\beta^2 Q_{3,0} - \beta(\gamma\beta + 2\alpha\delta) Q_{2,1} + \alpha(2\gamma\beta + \alpha\delta) Q_{1,2} - \alpha^2\gamma Q_{0,3}\right), \\
\bar{Q}_{0,3} &= \frac{1}{A^3}\left(-\beta^3 Q_{3,0} + 3\alpha\beta^2 Q_{2,1} - 3\alpha^2\beta Q_{1,2} + \alpha^3 Q_{0,3}\right),
\end{aligned}$$

where, as above, $A = \delta\alpha - \gamma\beta$. Equations

$$\Phi_{2,0} = 0,\ \Phi_{0,2} = 0,\ \Phi_{3,0} = 1,\ \Phi_{0,3} = 1.$$

are equivalent to

$$\bar{Q}_{2,0} = 0,\ \bar{Q}_{0,2} = 0,\ \bar{Q}_{3,0} = 1,\ \bar{Q}_{0,3} = 1. \tag{16}$$

From the first pair of equations (16) we deduce that the ratios $\frac{\delta}{\gamma}$ and $\frac{\beta}{\alpha}$ are two roots of the same quadratic equation $\hat{Q}_{2,0} Z^2 - 2\hat{Q}_{1,1} Z + \hat{Q}_{0,2}$. The discriminant of this quadratic equation is

$$d = \hat{Q}_{1,1}^2 - \hat{Q}_{2,0}\,\hat{Q}_{0,2} = \frac{1}{n} u H \tag{17}$$

where

$$H = n\,u(u_{pq}^2 - u_{pp}\,u_{qq}) + (n-1)\,(u_{pp}\,u_q^2 + 2\,u_p\,u_q\,u_{pq} + u_{qq} u_p^2) \tag{18}$$

is the inhomogenization of the Hessian of the polynomial $u = F(x, y, z)$. The roots are:

$$\frac{\delta}{\gamma} = r_1 = \frac{\hat{Q}_{1,1} + \sqrt{d}}{\hat{Q}_{2,0}}, \qquad \frac{\beta}{\alpha} = r_2 = \frac{\hat{Q}_{1,1} - \sqrt{d}}{\hat{Q}_{2,0}} \tag{19}$$

By subtracting these expressions one obtains that

$$r_1 - r_2 = \frac{\alpha\delta - \beta\gamma}{\alpha\gamma} = \frac{2\sqrt{d}}{\hat{Q}_{2,0}} \Longrightarrow\ A = \alpha\gamma\,\frac{2\sqrt{d}}{\hat{Q}_{2,0}}.$$

From the second pair of equations (16) we obtain that:

$$\begin{aligned} \alpha &= \frac{\hat{Q}_{2,0}}{\sqrt{d}} \left(r_1^3 Q_{3,0} - 3 r_1^2 Q_{2,1} + 3 r_1 Q_{1,2} - Q_{0,3} \right)^{1/3}, \\ \gamma &= \frac{\hat{Q}_{2,0}}{\sqrt{d}} \left(-r_2^3 Q_{3,0} + 3 r_2^2 Q_{2,1} - 3 r_2 Q_{1,2} + Q_{0,3} \right)^{1/3}. \end{aligned} \tag{20}$$

Since $\beta = \alpha\, r_2$, $\delta = \gamma\, r_1$ then:

$$\begin{aligned} \beta &= \frac{\hat{Q}_{1,1} - \sqrt{d}}{\sqrt{d}} \left(r_1^3 Q_{3,0} - 3 r_1^2 Q_{2,1} + 3 r_1 Q_{1,2} - Q_{0,3} \right)^{1/3}, \\ \delta &= \frac{\hat{Q}_{1,1} + \sqrt{d}}{\sqrt{d}} \left(-r_2^3 Q_{3,0} + 3 r_2^2 Q_{2,1} - 3 r_2 Q_{1,2} + Q_{0,3} \right)^{1/3}. \end{aligned} \tag{21}$$

The substitution of the group parameters (20, 21) into $\bar{Q}_{1,1}$, $\bar{Q}_{2,1}$ and $\bar{Q}_{1,2}$ produces three *fundamental differential invariants* $I_{1,1}, I_{2,1}, I_{1,2}$ of the action (9). This means that these functions, defined on J^3, are unchanged under prolonged action (9, 11) and, moreover, any other invariant function can be locally expressed in terms of $I_{1,1}, I_{2,1}, I_{1,2}$. Functions $I_{1,1}, I_{2,1}, I_{1,2}$ are independent and their restrictions to $\mathcal{K}$ are just the standard coordinate functions: $I_{1,1}|_{\mathcal{K}} = u_{1,1}$, $I_{2,1}|_{\mathcal{K}} = u_{2,1}$, $I_{1,2}|_{\mathcal{K}} = u_{1,2}$. Evaluation of functions $I_{1,1}, I_{2,1}, I_{1,2}$ at a point $\mathbf{u}^{(3)} \in J^3$ produces the image under the projection π of $\mathbf{u}^{(3)}$ to $\mathcal{K}$ along its orbit. Thus one can compute the signature manifold of a cubic $f(p,q)$ by evaluation $I_{1,1}, I_{2,1}, I_{1,2}$ at f, that is, by substitution of $u_{i,j}$ with the actual derivatives $\frac{\partial^{i+j} u}{\partial^i p \partial^j q}$. As the result we obtain three functions in p and q which define the signature manifold $\mathcal{C}_f$ parametrically. Since two different parameterizations can define the same manifold, in order to compare the signatures of two different cubics f and $\bar{f}$, we need to eliminate p and q and compare the corresponding implicit equations $\Psi_1\left(I_{1,1}|_f, I_{2,1}|_f, I_{1,2}|_f\right) = 0$ and $\Psi_2\left(I_{1,1}|_{\bar{f}}, I_{2,1}|_{\bar{f}}, I_{1,2}|_{\bar{f}}\right) = 0$ (see Remark 2.2 below). The difficulty is that the formulae for $I_{1,1}, I_{2,1}, I_{1,2}$ involve radicals and thus are unsuitable for polynomial elimination algorithms. They can be used, however, to build three independent rational invariants that can serve the same purpose. The most immediate choice consists of $I_{1,1}^3$, $I_{1,2} I_{2,1}$ and $I_{2,1}^3 + I_{1,2}^3$. Unfortunately these invariants have quite complicated formulae in terms of the coordinates on J^3. By applying some ideas of the classical invariant theory we have obtained simpler invariants

$$\begin{aligned} I_1 &= \frac{I_{2,1}\, I_{1,2} - 1}{{I_{1,1}}^3}, \qquad I_2 = -4\, \frac{1 + 9\, I_{2,1}\, I_{1,2}}{{I_{1,1}}^3}, \\ I_3 &= \frac{1 - 3\, {I_{2,1}}^2\, {I_{1,2}}^2 - 6\, I_{2,1}\, I_{1,2} + 4\, {I_{1,2}}^3 + 4\, {I_{2,1}}^3}{{I_{1,1}}^6}. \end{aligned}$$

It follows from Remark 2.3 below that $I_{1,1}|_{\bar{f}}$ is zero if and only if the ternary form f can be transformed into a binary form. From now on let us assume that

$I_{1,1}|_{\bar{f}} \neq 0$ holds. The restriction of I_1, I_2, I_3 to f gives a parametric description of its signature manifold $\mathcal{C}_f$. We discuss now how to compute the corresponding implicit equations, characterizing the smallest variety containing $\mathcal{C}_f$.

We assume that none of restriction $I_1|_f$, $I_2|_f$, $I_3|_f$ are identically zero. (The following discussion is easy to adapt otherwise.) By clearing denominators, the relations defining $I_1|_f$, $I_2|_f$, $I_3|_f$ lead to a set Σ of 3 polynomials $D_k\, I_k|_f - N_k$ (for $k = 1 \cdots 3$) in the polynomial ring $\mathbb{C}[I_1|_f, I_2|_f, I_3|_f, p, q]$ where D_k, N_k are polynomials in $\mathbb{C}[p, q]$ and such that $D_k \neq 0$ holds. Clearly Σ is a regular chain (see Section 2 in [Boulier et al., 2001b]) in $\mathbb{C}[I_1|_f, I_2|_f, I_3|_f, p, q]$ w.r.t. the variable ordering $I_1|_f > I_2|_f > I_3|_f > p > q$. Observe that the signature manifold $\mathcal{C}_f$ is contained in the set of regular zeros of Σ over $\mathbb{C}$. See [Aubry et al., 1999] for these notions related to regular chains. Let $\mathcal{J}$ be the saturated ideal of Σ. Since Σ is a regular chain, it is also a characteristic set of $\mathcal{J}$ w.r.t $I_1|_f > I_2|_f > I_3|_f > p > q$.

Observe now that $\mathcal{J}$ is a prime ideal. So, let T be the regular chain which is a characteristic set of $\mathcal{J}$ w.r.t $p > q > I_1|_f > I_2|_f > I_3|_f$ computed by means of the PARDI algorithm [Boulier et al., 2001a]. Let R be the regular chain $T \cap \mathbb{C}[I_1|_f, I_2|_f, I_3|_f]$. Relation (2) in [Moreno Maza, 1999] shows that the saturated ideal of R is the elimination ideal $\mathcal{J} \cap \mathbb{C}[I_1|_f, I_2|_f, I_3|_f]$. Observe that $R \neq \emptyset$ holds. Then, Th .2 p. 130 in [Cox et al., 1997] shows that the variety associated with $\mathcal{J} \cap \mathbb{C}[I_1|_f, I_2|_f, I_3|_f]$ is the smallest variety containing $\mathcal{C}_f$. Finally, we have proved that R characterizes the smallest variety containing $\mathcal{C}_f$.

REMARK 2.2: *Let us call the smallest variety containing $\mathcal{C}_f$ the* signature variety *V_f of f. The signature manifold $\mathcal{C}_f$ may not fill the entire signature variety V_f neither in complex nor and real case. Over $\mathbb{C}$, however, there exist varieties $U_1, \ldots, U_\ell$ with dimension smaller than that of V_f such that $V_f = \mathcal{C}_f \cup U_1 \cup \cdots \cup U_\ell$. See for instance [Aubry et al., 1999]. It follows that distinct signature varieties correspond to distinct signature manifolds. This is not the case over the real numbers: two different signature manifolds $\mathcal{C}_{f_1}$ and $\mathcal{C}_{f_2}$ can lie in the same signature variety. See Section (6) of this paper and Example 8.69 in [Olver, 1999].*

Using the ALDOR implementation (by the second author) of the PARDI algorithm we found the signature manifolds for all canonical cubics in terms of the above invariants. These computations suggested a new set of invariants:

$$\begin{aligned} i_1 &= 10\,(6\,I_1 + 1), \\ i_2 &= 6\,I_2 + 126\,I_1 + 45\,I_3 - 10, \\ i_3 &= 10\,(9\,I_3 + 2). \end{aligned} \tag{22}$$

leading to exceptionally simple results listed in Section 3.

REMARK 2.3: *None of the invariants is defined when $I_{1,1} \equiv 0$ (or equivalently $d \equiv 0$). This happens if and only if the inhomogenization of the Hessian (18) is identically zero, and hence if and only if the ternary form can be transformed into a binary form (see [Olver, 1999] p234 for the remarks's on Hesse's "theorem").*

3. Canonical Forms.

In order to match a cubic with its class we need to compute the signatures for each of the canonical forms listed in Theorems 1.1 and 1.2.

EXAMPLE 3.1: *To compute the signature for $F = z(x^2 + y^2 + z^2)$ we first write its inhomogeneous version $f = p^2 + q^2 + 1$ and restrict invariants i_1, i_2 and i_3 to f. We obtain:*

$$\begin{aligned} i_1|_f &= 90\,\frac{(p^2+q^2+1)^2}{(p^2-3+q^2)^2}, \\ i_2|_f &= 270\frac{(p^2+q^2+1)^3}{(p^2-3+q^2)^3}, \\ i_3|_f &= 180\,\frac{(p^2+q^2+1)\,((p^2+q^2+3)^2-12)}{(p^2-3+q^2)^3} \end{aligned}$$

Elimination of p and q leads to the defining equations for the one-dimensional signature variety:

$$i_1\,(i_3 - i_2) + 30\,i_2 = 0, \quad 10\,i_2^2 - i_1^3 = 0.$$

A cubic $\bar{F}$ can be transformed to F by a complex linear change of variables if and only if $i_1|_{\bar{f}}, i_2|_{\bar{f}}, i_3|_{\bar{f}}$, define a one-dimensional variety and satisfy the above equations.

We list below the signature varieties for the canonical cubics. In order to match an arbitrary cubic f with its canonical form we need to restrict the three invariants i_1, i_2 and i_3 to f. In order do determine its class, one does not need to compute the signature of f itself, but just to check which of the following relations are satisfied. The latter computations are not difficult and were programed in MAPLE and ALDOR.

We recall also that the signature of f is the projection along the orbits of its prolonged graph to the cross-section $\mathcal{K}$. Since the graph of f is two-dimensional the absolute majority of cubics have two-dimensional signatures. The graphs of some cubics, however, are not transversal to the orbits, and thus their signatures are one- or zero-dimensional. According to Theorem 2.2 this indicates the increase in the dimension of the symmetry groups G_F. We will list the symmetry groups below, postponing details until Section 4.

THEOREM 3.1: *Every homogeneous cubic $F(x, y, z)$ can be transformed by a linear change of variables to one and only one of the canonical forms listed below. In order to determine its canonical form, one need to compute $i_1|_f, i_2|_f, i_3|_f$ and to determine which of the relations listed below they satisfy. The symmetry group of F is isomorphic to the symmetry group of its canonical form under conjugation (3).*

Irreducible cubics.

- Regular (Elliptic Curves):

  (1) $F \sim x^3 + \mu\, xz^2 + z^3 - y^2 z, \quad f \sim p^3 + \mu\, p + 1 - q^2,$
  non-equivalent for different values of μ^3;
  $a \neq 0$ (else $F \sim (3)$), $\mu^3 \neq -27/4$ (else $F \sim (5)$),
  $|G_F| = 18 \times 3,$
  $\boxed{27\, i_1^3 + 40\, \mu^3\, i_2^2 = 0.}$

  (2) $F \sim x^3 + xz^2 - y^2 z, \quad f \sim p^3 + p - q^2,$
  $|G_F| = 36 \times 3,$
  $\boxed{i_2 = 0.}$

  (3) $F \sim (x^3 + z^3 - y^2 z) \sim (x^3 + y^3 + z^3) \quad f \sim p^3 + 1 - q^2,$
  $|G_F| = 54 \times 3,$
  $\boxed{i_1 = 0.}$

- Singular:

  (4) $F \sim x^3 - y^2 z, \quad f \sim p^3 - q^2,$
  $G_F \sim$ 1-dimensional: $x \to x$, $y \to \alpha y$, $z \to \alpha^{-2} z,$
  $\boxed{i_1 = 0, \quad i_2 = 0.}$

  (5) $F \sim x^2(x + z) - y^2 z, \quad f \sim p^2(p + 1) - q^2$
  $|G_F| = 6 \times 3,$
  $\boxed{i_1^3 - 10\, i_2^2 = 0.}$

Reducible cubics:

- a linear and an irreducible quadratic factor:

  (6) $F \sim z(x^2 + yz), \quad f \sim (p^2 + q)$
  $G_F \sim$ 2-dimensional noncommutative (affine) group, generated by:
  $x \to x + \alpha z$, $y \to -2\alpha x + y - \alpha^2 z$, $z \to z,$
  $x \to \beta x$, $y \to \beta^4 y$, $z \to \beta^{-2} z,$
  $\boxed{i_1 = 0,\ i_2 = 0,\ i_3 = 0.}$

  (7) $F \sim z(x^2 + y^2 + z^2), \quad f \sim p^2 + q^2 + 1$
  $G_F \sim$ 1-dimensional (rotation in the xy) plane,
  $\boxed{i_1(i_3 - i_2) + 30\, i_2 = 0,\ 10\, i_2^2 - i_1^3 = 0.}$

- three linear factors:

  (8) non-coplaner $\Longleftrightarrow F \sim xyz, \quad f \sim p\, q;$
  $G_F \sim$ 2-dimensional:
  $x \to \alpha x$, $y \to \beta y$, $z \to \frac{1}{\alpha\beta} z.$
  $\boxed{i_1 = 90,\ i_2 = 270,\ i_3 = 180.}$

  (9) different, but coplaner $\Leftrightarrow F \sim x\, y\, (x + y), \quad f \sim p\, q(p + q)$
  $G_F \sim$ 3-dimensional direct product of arbitrary linear transformations

$z \mapsto \alpha x + \beta y + \gamma z$ and a finite subgroup of order 18 of linear transformations on the (x, y)-plane, which preserve $xy(x + y)$.

(10) two repeated $\Leftrightarrow F \sim x^2\, y, \quad f \sim p^2 q$
$G_F \sim$ 4-dimensional group of matrices generated by the transformations
$x \to \alpha\, x, \quad y \to \frac{1}{\alpha^2} y, \quad z \to \beta\, x + \gamma\, y + \delta\, z.$

(11) three repeated $\Leftrightarrow F \sim x^3, \quad f \sim p^3.$
$G_F \sim$ 4-dimensional direct product of $GL(2, \mathbb{C})$ and cyclic group Z_3 (linear transformations on the variables y and z and multiplication of x by a cubic root of 1).

REMARK 3.1: *The first class is actually a one-parametric family of equivalence classes. The classes of* $x^3 + \mu_1 xz^2 + z^3 - y^2 z$ *and* $x^3 + \mu_2 xz^2 + z^3 - y^2 z$ *are different unless* $\mu_1^3 = \mu_2^3$. *In the latter case the transformation* $x \to \omega x$, *where* $w = \mu_1/\mu_2$ *is a cubic root of 1, maps the first cubic to the second one.*

REMARK 3.2: *The last three classes (9), (10) and (11), in fact, depend on less than three variables. Invariants* i_1, i_2, i_3 *are not defined for such polynomials, neither the procedures for computing symmetry groups and linear transformations described in the next two section. This case should be studied with the invariants applicable for binary forms (see [Olver, 1999], [Berchenko and Olver, 2000]).*

The picture below shows the signatures of canonical cubics in the three-dimensional space parameterized by i_1, i_2, i_3.

We notice that all low dimensional signatures are contained in some signatures of the higher dimension. This have a nice geometrical interpretation. As it has been mentioned in the introduction, the set of ternary homogeneous cubics form a ten-dimensional linear space W isomorphic to the symmetric tensor product $S^3\mathbb{C}^3$. The action (2) defines an irreducible representation on W. An inclusion of the signature of f into the signature of $\bar{f}$ indicates that the closure of the orbit of $\bar{f}$ contains the orbit of f.

EXAMPLE 3.2: *A cubic* $x^3 + \mu\, xz^2 + z^3 - y^2 z$ *from the one parametric family of equivalence classes (1) is transformed to* $x^3 + \mu\, \varepsilon^4 xz^2 + \varepsilon^6\, z^3 - y^2 z$ *under the linear map*

$$x \to x, \quad y \to \frac{1}{\varepsilon} y, \quad z \to \varepsilon^2 z. \tag{23}$$

The latter cubic tends to the cubic $x^3 - y^2 z$ *of class (4) when* $\varepsilon \to 0$. *The signature of cubics of class (4), the* i_3*-axis, is included into the signature of* $x^3 + \mu\, xz^2 + z^3 - y^2 z$ *for all* μ.

We summarize the inclusions of the signatures, and therefore the inclusions of one orbits in the closure of another, by the diagram shown on Figure (3). Note that inclusions (9), (10), (11) of binary forms are not seen from the signatures.



Figure 2: Signatures of canonical cubics.



Figure 3: Inclusions of signature manifolds.

The above picture can be compared with the analysis of the closures of the orbits under the action of the *special linear group* $SL(3,\mathbb{C})$ in Kraft [1985].

The linear space of cubics is ten-dimensional, while the acting group $GL(3,\mathbb{C})$ is nine-dimensional, and thus we expect to have one invariant for this action. This invariant will be an absolute invariant in the sense of the classical invariant theory. This means that it depends on the coefficients of a cubic, not on the variables p, q . We notice that the ratio i_1^3/i_2^2 provides such invariant.

PROPOSITION 3.1: *The ratio* $A = -\frac{27}{40}\frac{i_1^3}{i_2^2}$ *is an absolute invarinat for the action (2), that is, it depends only on the coefficients of a cubic, and is constant for each of the equivalence classes, whenever it is defined.*

Observe that A distinguishes equivalence classes in the set of cubics whose orbits have maximal dimension. Iit does not distinguish the equivalence classes in the set of all cubics. For instance, it equals to $-\frac{27}{4}$ for classes (5), (7) and (8).

4. The symmetry groups.

According to Definition 1.2 the *symmetry group* of a cubic F is the subgroup $G_F \subset GL(3,\mathbb{C})$ consisting of all linear transformations that map F to itself. It coincides with the group $G_f \subset GL(3,\mathbb{C})$ which maps inhomogeneous version f of F to itself under transformation (9). Due to Theorem 2.2 the dimension of the signature $\mathcal{C}_f$ determines the dimension of the symmetry group: $\dim G_f = 2 - \dim \mathcal{C}_f$. Since it is useful for computations of the symmetry groups, we sketch the main ideas underlining the proof of Theorem 2.2. We start with the following simple proposition:

PROPOSITION 4.1: *Assume* $\mathbf{u}^{(3)}$ *and* $\bar{\mathbf{u}}^{(3)}$ *are two points on the prolonged graph* $\Gamma_f^{(3)} \subset J^3$*, which lie on the same orbit. That means there exists a matrix* $g \in GL(3,\mathbb{C})$*, such that* $g \cdot \mathbf{u}^{(3)} = \bar{\mathbf{u}}^{(3)}$*. Then* $g \in G_f$*.*

Proof. The graphs $\Gamma_f^{(3)}$ and $g \cdot \Gamma_f^{(3)}$ are two prolonged graphs that pass through the point $\bar{\mathbf{u}}^{(3)}$, but there is only one such graph through each point. Thus $g \cdot \Gamma_f^{(3)} = \Gamma_f^{(3)}$, and hence $f = g \cdot f$. □

It follows that in order to find G_F one needs to study the intersection of the prolonged graph $\Gamma_f^{(3)}$ with the orbit $\mathcal{O}_{\mathbf{u}}^{(3)}$ of $\mathbf{u}^{(3)}$. In fact, since the isotropy group of each point of J^3 consists of three group elements, the points of the intersection $\Gamma_f^{(3)} \cap \mathcal{O}_{\mathbf{u}}^{(3)}$ are in one-to-three correspondence with the elements of the symmetry group. In order to determine this intersection, we recall that invariants i_1, i_2, i_3 define a projection π of J^3 onto the cross-section $\mathcal{K}$. Their restriction $i_1|_f, i_2|_f, i_3|_f$ to $\Gamma_f^{(3)}$ define a projection $\pi|_f \colon \Gamma_f^{(3)} \to \mathcal{K}$, whose image is the signature manifold of f.

DEFINITION 4.1: *A pair of coordinates* (p, q) *is called a generic point of a cubic* f*, if the corresponding point* $\mathbf{u}^{(3)}$ *on* $\Gamma_f^{(3)}$*, computed by formulae (10)*

1) does not annihilate d (equivalently, (p,q) neither lies in the projective variety defined by f, nor annihilates the Hessian of f);

2) projects to a generic point c on the signature manifold $\mathcal{C}_f$. That means that the point c is a non-singular point of the variety $\mathcal{C}_f$ and it does not belong to any lower-dimensional signature manifold.

A randomly chosen point will be generic with probability one. Let us choose a generic point (p_0, q_0), and compute the corresponding point $\mathbf{u}^{(3)} \in \Gamma_f^{(3)}$ and its projection $c = \pi|_f(\mathbf{u}^{(3)}) \in \mathcal{C}_f$. Since $\pi|_f$ defines the projection along the orbits, then the preimage $\pi|_f^{-1}(c) = \Gamma_f^{(3)} \cap \mathcal{O}_{\mathbf{u}^{(3)}}$ is the desired intersection. Moreover,

$$\dim G_f = \dim \pi|_f^{-1}(c) = \dim \Gamma_f^{(3)} - \dim \mathcal{C}_f = 2 - \dim \mathcal{C}_f.$$

If the signature manifold has a non maximal dimension, then a cubic has a continuous group of symmetries of dimension one or two. The corresponding linear transformations, listed in Theorem 3.1, can be easily found using Lie's infinitesimal methods [Olver, 1986], proposition 2.6. When a signature manifold of f has the maximal dimension, two, the corresponding cubic has a finite group of symmetries which can be explicitly found by solving the equations:

$$\begin{aligned} i_1|_f(p,q) = i_1|_f(P,Q), \qquad & i_2|_f(p,q) = i_2|_f(P,Q), \\ i_3|_f(p,q) \;=\; & i_3|_f(P,Q) \end{aligned} \tag{24}$$

for P and Q in terms of p and q. Indeed, the solution of this system will determine all points on Γ_f which are projected under π_f to the same point on the signature $\mathcal{C}_f$. Remarkably all the solutions of the equations (24) are linear fractional (see [Olver, 1999] chapter 8, p. 190 and [Berchenko and Olver, 2000]):

$$\begin{aligned} P &= \frac{\alpha\, p + \beta\, q + \lambda}{a\, p + b\, q + \eta}, \\ Q &= \frac{\gamma\, p + \delta\, q + \mu}{a\, p + b\, q + \eta}. \end{aligned}$$

These symmetries are called *projective*. Each projective symmetry gives rise to three genuine symmetries of the form $f(p,q)$. In practice however the equations (24) are difficult to solve. Nevertheless, fixing a specific generic point (p_0, q_0), one can find the number of the solutions using a well known algebraic result ([Cox et al., 1997], Proposition 8, ch. 5, § 3) and the Gröbner basis computation (see [Cox et al., 1998], ch. 2,§ 2). The number of the solutions determines the cardinality of the symmetry group of f. We conclude this section with a simple corollary from Theorems 3.1 and formula (3).

COROLLARY 4.1: *A homogeneous cubic F in three variables is irreducible if and only if $G_F \cap SL(3,\mathbb{C})$ is discrete. A cubic F is a product of a linear factor and an irreducible quadratic factor if and only if its $G_F \cap SL(3,\mathbb{C})$ is one-dimensional. A cubic F is a product of three different linear factors if and only if its $G_F \cap SL(3,\mathbb{C})$ is two-dimensional.*

5. Transforming a Cubic to its Canonical Form.

Given an arbitrary cubic polynomial F we can match it with one of the canonical forms listed in Theorem 3.1 by computing invariants $i_1|_f, i_2|_f, i_3|_f$ and determining which of the relations listed in Theorem 3.1 they satisfy. From Theorem 2.1 we know that F can be mapped to the corresponding canonical form $\bar{F}$ by a linear change of variables g. We would like to compute g explicitly. We first note that we should not expect a unique answer. Indeed, if $\bar{F} = g \cdot F$ and $h \in G_F$, then also $\bar{F} = gh \cdot F$. In fact, the entire left coset gG_F will map F to $\bar{F}$. Thus the number of possible transformations depends on the size of the symmetry group of F. In the finite case one can find all such transformations by solving the system of equations:

$$\begin{aligned} i_1|_f(p,q) = i_1|_{\bar{f}}(P,Q), \qquad & i_2|_f(p,q) = i_2|_{\bar{f}}(P,Q), \\ i_3|_f(p,q) \;\; = \;\; & i_3|_{\bar{f}}(P,Q) \end{aligned} \tag{25}$$

for P, Q in terms of p, q, where f and $\bar{f}$ are inhomogeneous versions F and $\bar{F}$ respectively (see [Olver, 1999] Chapter 8). This computation, however, is too difficult to be performed in practice and non-applicable in the infinite case, therefore, we propose an alternative more practical algorithm.

Given an inhomogeneous version f of F one can compute the elements of the matrix g (see (4)) using formulae (14, 20, 21). The entries of the obtained matrix are some functions of (p, q) and therefore we denote the matrix by $g|_f(p, q)$. Let us choose specific numeric values (p_0, q_0) that are generic in the sense of Definition 4.1 and denote a corresponding matrix with constant entries as g. By construction, g brings $\mathbf{u}^{(3)}$ to a point on the signature $\mathcal{C}_f$: $g \cdot \mathbf{u}^{(3)} = c \in \mathcal{C}_f$. Let $\bar{f}$ be equivalent to f, then $\mathcal{C}_{\bar{f}} = \mathcal{C}_f$, and so we can find a point (P_0, Q_0) such that the corresponding point $\bar{\mathbf{u}}^{(3)} \in \Gamma^{(3)}_{\bar{f}}$ is projected to the same point c on the signature manifold. This can be done by solving for P, Q the system (25) with $p = p_0, q = q_0$. Although the system we consider has many, or even infinitely many solutions, depending on the size of the symmetry group G_f, we are interested in any one of the solutions (P_0, Q_0). The corresponding point $\bar{\mathbf{u}}^{(3)} \in \Gamma^{(3)}_{\bar{f}}$ is projected to the same point $c \in \mathcal{C}_f$ as the point $\mathbf{u}^{(3)}$. Using again formulae (14, 20, and 21), we compute the matrix $\bar{g}|_{\bar{f}}(p, q)$, that corresponds to $\bar{f}$ and evaluate it at the point (P_0, Q_0) to obtain the matrix $\bar{g}$. The matrix $\bar{g}^{-1}\, g$ is the desired linear transformation.

This procedure can be summarized as follows.

Step 1 Given a homogeneous cubic $F(x, y, z)$ compute its inhomogeneous version $f(p, q) = F(p, q, 1)$ and the corresponding differential invariants given by relation $i_1|_f, i_2|_f, i_3|_f$ (22).

Step 2 Determine the canonical form $\bar{F}$, by determining which of the relations listed in Theorem 3.1 are satisfied and compute the corresponding invariants $i_1|_{\bar{f}}, i_2|_{\bar{f}}, i_3|_{\bar{f}}$.

STEP 3 Choose a generic point (p_0, q_0) of f and find a corresponding point (P_0, Q_0) for $\bar{f}$ as a solution of equation (25).

STEP 4 Compute matrices $g = g|_f(p_0, q_0)$ and $\bar{g} = g_{\bar{f}}(P_0, Q_0)$ using (14, 20, and 21).

STEP 5 The matrix $g_0 = \bar{g}^{-1}\, g$ is the desired linear transformation, such that we have $\bar{F}(g_0 \cdot \mathbf{x}) = F(\mathbf{x})$.

The implementation of STEP 2 and STEP 3 requires some comments, meanwhile the other steps are straightforward.

Because of the inclusions of the signatures shown on Figure (3), performing STEP 2 requires to determine $\bar{F}$ by testing the membership of F to each of the first 8 classes of Theorem 3.1 sorted by increasing dimension of their signature manifolds. For instance: (6), (8), (4), (7), (2), (3), (5) and (1).

Performing STEP 3 implies solving a system consisting of non linear equations. This task can be achieved very easily in each of the first 8 cases of Theorem 3.1. Indeed, as we shall see now, in the worst case, the most expensive operations are

(a) the computation of a resultant of two bivariate polynomials with rational coefficients and with total degree not greater than 11,

(b) the factorization of this resultant,

(c) the computation of the gcd of the univariate polynomials obtained by specializing the initial system at each root of the resultant.

In practice we may want to restrict this process to the computation of a single solution (P_0, Q_0) which may speed up the above steps (b) and (c).

The method described in [Bix, 1998] (for computing a linear transformation that maps a ternary cubic to its canonical form) requires similar operations: solving systems of bivariate equations, factorization. We detail now the computations of our STEP 3 in each of the 8 cases of Theorem 3.1.

For cases (6) and (8) the invariants $i_1|_f$, $i_2|_f$ and $i_3|_f$ are constant and thus we can choose any point (P_0, Q_0). For the cases (1), (2), and (5) the quantities $i_1|_{\overline{f}}$, $i_2|_{\overline{f}}$ and $i_3|_{\overline{f}}$ are given by

$$i_1|_{\overline{f}} = k_1 \frac{\overline{f}^2}{\overline{V}^2}, \quad i_2|_{\overline{f}} = k_2\, \frac{\overline{f}^3}{\overline{V}^3} \quad \text{and} \quad i_3|_{\overline{f}} = k_3 \frac{\overline{f}\,\overline{W}}{\overline{V}^3}$$

where k_1, k_2, k_3 are constant and where $\overline{V}$, $\overline{W}$ are polynomial functions of (p, q) with total degree 3 and 6 respectively. Let us denote by $\overline{V}_1$, $\overline{V}_2$, $\overline{V}_5$ and $\overline{W}_1$, $\overline{W}_2$, $\overline{W}_5$ the corresponding values of $\overline{V}$ and $\overline{W}$ in case (1), (2), and (5) respectively.

One can easily check that:

$$\begin{aligned}
\overline{V}_1 &= \mu^2 - 9\,p - 3\,p\,q^2 - 3\,\mu\,p^2 \\
\overline{W}_1 &= -2\,\mu^4\,p + 4\,\mu^3\,p^3 + \mu^3 + 7\,\mu^3\,q^2 + 18\,\mu^2\,p^5 + 30\,\mu^2\,p^2\,q^2 - 9\,\mu^2\,p^2 - 9\,p^4\,q^2\,\mu \\
&\quad +27\,p^4\,\mu + 54\,\mu\,p\,q^2 - 6\,\mu\,p\,q^4 - 27\,p^6 + 54\,p^3 - 54\,p^3\,q^2 + q^6 - 18\,q^4 + 81\,q^2 \\
\overline{V}_2 &= 3\,p\,q^2 - 1 + 3\,p^2 \\
\overline{W}_2 &= 18\,p^5 - 9\,q^2\,p^4 + 4\,p^3 + 30\,p^2\,q^2 - 6\,p\,q^4 + 2\,p + 7\,q^2 + q^6 \\
\overline{V}_5 &= -3\,p\,q^2 - q^2 + p^2 \\
\overline{W}_5 &= 6\,p^6 + 6\,p^5 + 2\,p^4 - 9\,p^4\,q^2 - 4\,p^2\,q^2 - 6\,p\,q^4 - 3\,q^6 + 2\,q^4
\end{aligned}$$

The triple of constants (k_1, k_2, k_3) is $(-270\,\mu, 3645, -270)$ in case (1), $(-270, 0, 270)$ in case (2) and $(90, 270, 90)$.

Similarily the quantities $i_1|_f$, $i_2|_f$ and $i_3|_f$ are of the form:

$$i_1|_f = k_1 \frac{f^2}{V^2}, \quad i_2|_f = k_2\ \frac{f^3}{V^3} \quad \text{and} \quad i_3|_f = k_3 \frac{fW}{V^3}\,.$$

It follows that the system (25) simplifies as follows:

$$\frac{f}{V} = \frac{\overline{f}}{\overline{V}} \quad \text{and} \quad \frac{W}{V^2} = \frac{\overline{W}}{\overline{V}^2}\,. \tag{26}$$

In Step 3 the left hand sides of these equations are specialized at a numeric point (p_0, q_0). Hence the unknown point (P_0, Q_0) is one of the solutions of a system of total degree 6. Moreover one can check easily that this system has a finite number of complex solutions (which is of no surprise since the symmetry groups are finite in these cases). For case (3) of Theorem 3.1 the quantities $i_1|_{\overline{f}}$, $i_2|_{\overline{f}}$ and $i_3|_{\overline{f}}$ are given by

$$i_1|_{\overline{f}} = 0, \quad i_2|_{\overline{f}} = 135\ \frac{\overline{f}^3}{\overline{V}_3^3} \quad \text{and} \quad i_3|_{\overline{f}} = -10 \frac{\overline{f}\,\overline{W}_3}{\overline{V}_3^3} \tag{27}$$

where $\overline{V}_3 = p\,q^2 + 3\,p$ and $\overline{W}_3 = q^6 - 18\,q^4 + 81\,q^2 - 54\,p^3\,q^2 + 54\,p^3 - 27\,p^6$. The system (25) splits into two systems: one with total degree 6 and the other with total 12. Moreover both have a finite number of complex solutions.

For cases (4) and (7) the system (25) reduces to a single equation of the form

$$\frac{f}{V} = \frac{\overline{f}}{\overline{V}} \tag{28}$$

where $V_4 = p^3$ and $V_7 = p^2 - 3 + q^2$. Hence f, V have degree 3 in case (4) and degree 2 in case (7). To compute a point (P_0, Q_0) in this situation we could choose randomly one of its coordinate and then solve the corresponding univariate equation deduced from (28). However, we try to find rational values for P_0

and Q_0 in order to avoid the introduction of algebraic numbers. For instance, for case (4) we use the following remark. (This is certainly a classical observation.)

Let K be a commutative field with characteristic different from 2 and 3. Let $g(P) = P^3 + X\,P^2 + Y\,P + Z$ be a univariate polynomial in $K[P]$. The system of equations consisting of $g(P)$ and its derivative $g'(P) = 3\,P + 2\,X\,P + Y$ has the following triangular decomposition in $K[P, X, Y, Z]$ w.r.t. $P > X > Y > Z$:

$$
\left\{
\begin{array}{l}
(2X^2 - 6Y)\,P + YX - 9Z \\
4ZX^3 - Y^2X^2 - 18ZYX + 4Y^3 + 27Z^2
\end{array}
\right.
\left\{
\begin{array}{l}
3P + X \\
YX - 9Z \\
Y^3 - 27Z^2
\end{array}
\right.
\left\{
\begin{array}{l}
2P + X \\
X^2 - 4Y \\
Z
\end{array}
\right.
\left\{
\begin{array}{l}
P \\
Y \\
Z
\end{array}
\right.
$$

This shows that if $g(P)$ has a double root in some extension field of K then this root belongs in fact to K. Since X is equal to the sum of the root of g, it follows that if the discriminant of g vanishes then all the roots of g belong to K. Consider now the case where K is the field of fractions of $\mathbb{Q}[Q]$ and let G be a polynomial in $K[P, Q]$ such that $G = k\,g$ for some $k \in K$. The previous statement shows that if the numerator of the discriminant of g has a rational root then there exists a point (P_0, Q_0) with rational coordinates such that $G(P_0, Q_0) = 0$. Therefore, to solve the equation (28), say $G(P, Q) = 0$, we consider the additional equation given by $\Delta(Q) = 0$ where $\Delta(Q)$ is the discriminant of G w.r.t. Q. In practice, when there is a rational solution $G(P, Q) = 0$ then this *discriminant trick* always provides us with a rational point (P_0, Q_0). Indeed, it turns out that G is itself a ternary cubic of case (4) in inhomogeneous form and it is known that if such cubic has a rational point then its singular point is also rational [Shafarevich, 1994].

When solving the system (25) with a fixed point (p_0, q_0) we are finding the set of points (P_0, Q_0) which have the same image on the signature mantels as (p_0, q_0). Thus if (p_0, q_0) is generic then so is (P_0, Q_0), and conversely. We need now to explain how we do obtain a generic point. In our implementation of the algorithm we solve this question as follows. For each canonical form we have precomputed a generic point (P_0, Q_0). Then we solve the system (25) for a point (p_0, q_0). The equations to solve are similar to those of the converse way. In particular, for each case, we have the same upper bound for the total degree. However they are generally less sparse. This explains how we were led to the above discriminant trick.

6. Classification over $\mathbb{R}$.

Two polynomials f_1 and f_2 with real coefficients, that are equivalent under a complex linear change of variables, might not be equivalent under the real changes of variables. The simplest example is provided by polynomials p^2+q^2 and p^2-q^2. In fact, most of the complex equivalence classes splits into two real subclasses. Below we provide a real classification (irreducible case can be found in [Bix, 1998]) and discuss how to apply the moving frame method to match a real cubic with its real canonical form. We write $f_1 \sim_{\mathbb{C}} f_2$ if f_1 and f_2 are equivalent over complex transformations and $f_1 \sim_{\mathbb{R}} f_2$ if they are equivalent over real ones.

THEOREM 6.1: *A homogeneous cubic $f(p,q)$ with real coefficients and non-zero Hessian can be transformed by a real transformation (9) to one and only one of the canonical forms listed below.*

Since they are useful in determination of the real class, we will list the cubics along with a real point (p_0,q_0), generic in the sense of definition 4.1. Note, in particular, that this point *is not* on the variety defined by f. We also list the value of d *at this point.*

Irreducible cubics.

- Regular (Elliptic Curves):

  (1) (a) $(p^3+p+h-q^2) \sim_{\mathbb{C}} (p^3+h^{-2/3}p+1-q^2)$,
  $(p_0,q_0)=(0,1), d=\frac{4}{3}(1-h)$, (for $h=1$, $(p_0,q_0)=(1,0), d=44$).
  if $h>0$ then $(p^3+p+h-q^2) \sim_{\mathbb{R}} (p^3+h^{-2/3}p+1-q^2)$.

  (b) $(p^3+p-h-q^2) \sim_{\mathbb{C}} (p^3-h^{-2/3}\,p+1-q^2)$,
  $(p_0,q_0)=(0,1), d=\frac{4}{3}(1+h)$, (for $h=-1$, $(p_0,q_0)=(1,0), d=44$).
  if $h>0$ then $(p^3+p-h-q^2) \sim_{\mathbb{C}} (p^3-h^{-2/3}\,p+1-q^2)$

  In both cases cubics with different values of h are not equivalent over the reals. Two cubics that differ only by the sign of h have the same canonical form over complex numbers and are not equivalent over the real numbers. If $h=0$ then $f \sim (2)$ and if $h^2 \neq \frac{4}{27}$ then $f \sim (5)$).

  (2) (a) p^3+p-q^2, $(p_0,q_0)=(1,0), d=\frac{16}{3}$.
  (b) p^3-p-q^2, $(p_0,q_0)=(1,0), d=\frac{4}{3}$.

  (3) (a) p^3+1-q^2, $(p_0,q_0)=(1,0), d=24$.
  (b) p^3-1-q^2, $(p_0,q_0)=(1,0), d=8$.

- Singular:

  (4) $f \sim p^3-q^2$, $(p_0,q_0)=(2,1), d=56$.

  (5) (a) $p^2(p+1)+1-q^2$, $(p_0,q_0)=(2,1), d=44$.
  (b) $p^2(p-1)-1-q^2$, $(p_0,q_0)=(2,1), d=4$.

Reducible cubics:

- a linear and an irreducible quadratic factor:

(6) $(p^2+q) \quad (p_0,q_0)=(1,0), d=\frac{4}{3}$.
(7) (a) $p^2+q^2+1, \quad (p_0,q_0)=(1,0), d=16$.
(b) $p^2-q^2+1, \quad (p_0,q_0)=(1,0), d=\frac{16}{3}$.
(c) $p^2+q^2-1, \quad (p_0,q_0)=(1,1), d=\frac{20}{3}$.

- three linear factors:

(8) (a) $p^2-q^2 \sim_{\mathbb{R}} pq, \quad (p_0,q_0)=(1,0), d=-\frac{4}{3}$.
(b) $p^2+q^2 \quad (p_0,q_0)=(1,0), d=\frac{4}{3}$.

Given an arbitrary cubic we first identify which of the eight complex classes it belongs. Then in most cases we have to choose between two possibilities. In cases (4) and (6) there is only one possibility. In case (7) there are three possibilities. Assume we want to decide whether $f \sim_{\mathbb{R}} \bar{f}$, where $\bar{f}$ is one of the canonical forms listed above. First, we look for a *real* solution (P_0,Q_0) of the system (25), with the chosen generic point $p=p_0, q=q_0$. Moreover we require that d (defined by (17)) have the same sign: $\mathtt{sign}(d|_{\bar{f}}(p_0,q_0)) = \mathtt{sign}(d|_f(P_0,Q_0))$. Note that this sign is positive except case 8(a). If there is no such real solutions then we have a "wrong" canonical form. Once we find such solution, we construct the matrices $\bar{g}=g|_{\overline{f}}(p_0,q_0)$ and $g=g|_f(P_0,Q_0)$. Note that these matrices are real if $d>0$ (that is, for all cases except case 8 (a)). In all cases matrix $\bar{g}^{-1}\,g$ is the desired real transformation.

EXAMPLE 6.1: *Let us prove that polynomials* $f_1=p^2+q^2+1$ *and* $f_2=p^2+q^2-1$ *are not equivalent under real transformation (9). Equivalently cubic* $F_1=z(x^2+y^2+z^2)$ *can not be transformed to* $F_2=z(x^2+y^2-z^2)$ *under a linear real change of variables. These polynomials are listed above as canonical forms 7(a) and 7(c). Invariants* i_1,i_2,i_3 *for case (7) parameterize the signature variety* $i_1(i_3-i_2)+30\,i_2=0,\ 10\,i_2^2-i_1^3=0$. *Thus equations (25) reduce to one condition* $i_2|_{f_1}(p_0,q_0)=i_2|_{f_2}(P_0,Q_0)$, *where*

$$i_2|_{f_1}=270\frac{(p^2+q^2+1)^3}{(p^2-3+q^2)^3} \text{ and } i_2|_{f_2}=270\frac{(p^2+q^2-1)^3}{(p^2+3+q^2)^3}$$

Let us choose a real generic point $p_0=1, q_0=0$ *then our equation becomes*

$$\frac{(P_0^2+Q_0^2+1)}{(P_0^2-3+Q_0^2)}=1.$$

The latter does not have real solutions.

Although in, general, finding real solutions for a system of three equations and one inequality is a difficult problem, it is simple in each particular case, we consider. For instance in cases (6) and (8) the equations are empty, so we only need to check the sign of d. Case (6) does not split over reals and $d|_f=\frac{4}{3}f$. In case 8(a) $d|_f=-\frac{4}{3}f^2$ is negative for all real (p,q), while in case 8 (b) it is positive: $d|_f=\frac{4}{3}f^2$. Case (4) does not split into subcases. In the cases (1), (2),

(3) and (5) equations (25) have finite number of solutions and thus we can check the sign of d for each of them. In case (7) the situation is always similar to the one described in the above example. If two real cubics of complex case (7) are not equivalent over reals, equations (25) do not have real solutions.

We note that, as in the complex case, the real classification of the cubics is known. In the irreducible case one first finds a real inflection point and maps it to infinity. After this several quite simple operations are necessary Bix [1998]. For reducible cubics the classification can be also obtained in elementary way. Our goal was to confirm the efficiency of the moving frame method, which can be applied to polynomials of higher degree.

7. Conclusions.

In this paper we conduct a careful study of the equivalence classes of ternary cubics under general linear changes variables via moving frame method. The main ideas of such application appeared first in Olver [1999].

The new contribution of this paper is the computation of the signature manifolds for each of the equivalence classes and a practically feasible algorithm that matches a cubic with its canonical form, producing explicitly a required linear transformation. The implementation is made possible by triangular decomposition methods. We also make an interesting observation of the correspondence between the geometry of signature manifolds and the geometry of the orbits in the linear space of ternary cubics under the action of $GL(3, \mathbb{C})$.

There is no theoretical difficulty in applying the same methods to the polynomials of higher degree or higher number of variables. For ternary forms of *any* degree n the problem can be solved at the jet space of order $\min\{n, 6\}$ and the invariants for cubics can be reused with addition of new invariants of higher orders. The rational invariants of order four were computed in [Kogan, 2000]. However, the computations become more challenging and we are unaware of the complete classifications results for ternary forms of higher degrees.

The advantage of the moving frame method lies in its generality: it is applicable to any equivalence problems under an arbitrary finite-dimensional group of transformation. In particular, instead of the general linear group one can consider its subgroups, such as the special linear group or the orthogonal group. In this paper the dehomogenization of the cubic $F(x, y, z)$ leads us to the linear fractional action (8) on polynomials in two variables of degree less or equal to three. One can, more generally, consider nonzero weight transformation rules $f \to \bar{f}$:

$$\bar{f} = (\det g)^k \, (ap + bq + \eta)^3 \, f\left(\frac{\alpha p + \beta q + \lambda}{ap + bq + \eta}, \frac{\gamma p + \delta q + \mu}{ap + bq + \eta}\right).$$

The affine action is also of interest.

Cartan's method of equivalence was formulated in the category of smooth manifolds. Hence, its direct applicability is restricted to polynomials over complex

or real numbers. It would be a worthwhile and interesting project to reformulate the method of moving frames in the algebraic-geometry language, so that it can be applied to the problem of the equivalence and symmetry of algebraic varieties over fields of arbitrary characteristics.

Acknowledgements: The first author would like to acknowledge the support of the NSF VIGRE grant DMS 99-83403. The second author would like to thank the Ontario Research for Computer Algebra (University of Western Ontario, Canada) for supporting his work.

References

M. Ackerman and Hermann R. Hilbert's Invariant Theory Papers. *Math Sci Press, Brookline, Mass.*, 8, 1978.

P. Aubry, D. Lazard, and M. Moreno Maza. On the theories of triangular sets. *J. of Symb. Comput.*, 28(1-2):105–124, 1999.

I. A. Berchenko and P. J. Olver. Symmetries of polynomials. *J. of Symb. Comput.*, 29:485–514, 2000.

R Bix. *Conics and cubics. A concrete introduction to algebraic curves.* Springer-Verlag, 1998.

F. Boulier, F. Lemaire, and M. Moreno Maza. PARDI ! In *proceedings of ISSAC 2001*, pages 38–47, London, Ontario, 2001a. ACM Press.

F. Boulier, F. Lemaire, and M. Moreno Maza. Well known theorems on triangular systems. Technical Report LIFL 2001–08, Univ. of Lille I, LIFL, 2001b.

M. Bronstein, M. Moreno Maza, and S.M. Watt. *The Aldor User Guide.* Univ. of Western Ontario, 2001.

É. Cartan. La méthode du repère mobile, la théorie des groupes continus, et les espaces généralisés. *Exposés de Géométrie*, 5:Hermann, Paris, 1935.

P. Common and B. Mourrain. Decompositions of quantics in sums of powers of linear forms. *Signal Processing*, 53:93–107, 1996.

D. Cox, J. Little, and D. O'Shea. *Ideals, Varieties, and Algorithms.* Undergraduate Text in Mathematics. Springer-Verlag, New-York, 1997.

D. Cox, J. Little, and D. O'Shea. *Using Algebraic Geometry.* Graduate Text in Mathematics, 185. Springer-Verlag, New-York, 1998.

M. Fels and P. J. Olver. Moving Coframes. II. Regularization and Theoretical Foundations. *Acta Appl. Math.*, 55:127–208, 1999.

J. H. Grace and A. Young. *The Algebra of Invariants*. Cambridge Univ. Press, Cambridge, 1903.

M. L. Green. The moving frame, differential invariants and rigidity theorems for curves in homogeneous spaces. *Duke Math. Journal*, 45:735–779, 1978.

P. A. Griffiths. On Cartan's method of Lie groups as applied to uniqueness and existence questions in differential geometry. *Duke Math. Journal*, 41:775–814, 1974.

H. W. Guggenheimer. *Differential Geometry*. McGraw-Hill, New York, 1963.

G. Gurevich. *Foundations of the theory of algebraic invariants*. Noordhoff, 1964.

R. Howe. Remarks on classical invariant theory. *Trans. Amer. Math. Soc.*, 313: 539–570, 1989.

M. Kalkbrener. A generalized euclidean algorithm for computing triangular representations of algebraic varieties. *J. of Symb. Comput.*, 15:143–167, 1993.

A. W. Knapp. *Elliptic Curves*. Mathematical Notes. Princeton Univ. Press, Princeton, 1992.

I. A. Kogan. *Inductive Approach to Cartan's Moving Frames Method with Applications to Classical Invariant Theory*. PhD thesis, Univ. of Minnesota, 2000.

I. A. Kogan. Inductive Construction of Moving Frames. *Contemp. Math.*, 285: 157–170, 2001.

I.A. Kogan and M. Moreno Maza. Computation of canonical forms for ternary cubics. In Teo Mora, editor, *ISSAC 2002*, pages 151–160. ACM Press, July 2002.

Kh. Kraft. *Geometrische Methoden in der Invariantentheorie*. Friedr. Vieweg & Sohn, Braunschweig, 1985.

S. Lang. *Elliptic Functions*. Graduate Texts in Mathematics. Springer, New York, 1987.

M. Moreno Maza. On triangular decompositions of algebraic varieties. Technical Report TR 4/99, NAG Ltd, Oxford, UK, 1999. Presented at the MEGA-2000.

D. Mumford. *Geometric Invariant Theory*. 3rd ed. Springer, New York, 1994.

P. J. Olver. *Applications of Lie Groups to Differential Equations*. Springer, New York, 1986.

P. J. Olver. *Equivalence Invariants and Symmetry*. Cambridge Univ. Press, Cambridge, 1995.

P. J. Olver. *Classical Invariant Theory.* Cambridge Univ. Press, Cambridge, 1999.

B. Reznick. Sums of even powers of real linear forms. *Mem. Amer. Math. Soc.*, 96:1063–1073, 1992.

É. Schost. *Sur la résolution des systèmes polynomiaux à paramètres.* PhD thesis, École Polytechnique, Palaiseau, France, 2001.

I. R. Shafarevich. *Basic algebraic geometry. 1. Varieties in projective space.* Springer-Verlag, second edition, 1994. Translated from the 1988 Russian edition and with notes by Miles Reid.

E. B. Vinberg and V. L. Popov. *Invariant theory.* Algebraic geometry, 4, (Russian). Itogi Nauki i Tekhniki, Akad. Nauk SSSR, VINITI, Moscow, 1989.