

Computing the Limit Points of Quasi-componets of Regular Chains in Diemnsion One

Marc Moreno Maza

Joint work with Parisa Alvandi and Changbo Chen

University of Western Ontario, Canada

ZIB, Berlin, CASC 2013

September 9, 2013

Plan

- 1 The problem
- 2 Motivation
- 3 An introductory example (informal)
- 4 A more advanced example (informal)
- 5 Limit points and Puiseux expansions of an algebraic curve
- 6 Puiseux expansions of a regular chain and $\lim(W(T))$
- 7 Computation of $\lim(W(T))$
- 8 Experimentation
- 9 Demo
- 10 Conclusion

Plan

- 1 The problem
- 2 Motivation
- 3 An introductory example (informal)
- 4 A more advanced example (informal)
- 5 Limit points and Puiseux expansions of an algebraic curve
- 6 Puiseux expansions of a regular chain and $\lim(W(T))$
- 7 Computation of $\lim(W(T))$
- 8 Experimentation
- 9 Demo
- 10 Conclusion

Specification of the problem

Input

- Let $R \subset \mathbb{C}[X_1, \dots, X_s]$ be a regular chain.
- Let h_R be the product of initials of polynomials of R .
- Let $W(R)$ be the quasi-component of R , that is $V(R) \setminus V(h_R)$.

Output

The non-trivial limit points of $W(R)$, that is $\overline{W(R)}^Z \setminus W(R)$, denoted by $\lim(W(R))$.

Plan

- 1 The problem
- 2 **Motivation**
- 3 An introductory example (informal)
- 4 A more advanced example (informal)
- 5 Limit points and Puiseux expansions of an algebraic curve
- 6 Puiseux expansions of a regular chain and $\lim(W(T))$
- 7 Computation of $\lim(W(T))$
- 8 Experimentation
- 9 Demo
- 10 Conclusion

Motivation (I): the Ritt problem

The Ritt problem

Given the characteristic sets of two prime differential ideals $\mathcal{I}_1$ and $\mathcal{I}_2$, determine whether $\mathcal{I}_1 \subseteq \mathcal{I}_2$ holds or not:

- No algorithm is known,
- Equivalent to other key problems, see (O. Golubitsky et al., 2009).

The algebraic counterpart of the Ritt problem

Given regular chains R_1 and R_2 , determine whether $\text{sat}(R_1) \subseteq \text{sat}(R_2)$ holds or not, **without** computing a basis for $\text{sat}(R_1)$ or $\text{sat}(R_2)$:

- No algorithm is known,
- Such an algorithm could be used to solve the differential problem.

Our strategy for the algebraic version

- $\sqrt{\text{sat}(R_1)} \subseteq \sqrt{\text{sat}(R_2)} \iff \overline{W(R_2)}^Z \subseteq \overline{W(R_1)}^Z$
- $\overline{W(R)}^Z = W(R) \cup \text{lim}(W(R))$

Motivation (II): from Kalkbrener to Wu-Lazard decompositions

Specification (in the case of an irreducible variety)

Input: An irreducible algebraic set $V(F)$ and a regular chain R s.t.
$$V(F) = \overline{W(R)}^Z$$

Output: Regular chains $R_1, \dots, R_e$ s.t.
$$V(F) = W(R_1) \cup \dots \cup W(R_e)$$

Wu's trick

- Compute a triangular decomposition of $F \cup \{h_R\}$.
- The trick generalizes to the case where $V(F)$ is not irreducible.
- In practice, this process is very inefficient (many repeated calculations).

Our proposed strategy

- Compute $V(F) \setminus W(R)$ directly as the set $\text{lim}(W(R))$.

Plan

- 1 The problem
- 2 Motivation
- 3 An introductory example (informal)**
- 4 A more advanced example (informal)
- 5 Limit points and Puiseux expansions of an algebraic curve
- 6 Puiseux expansions of a regular chain and $\lim(W(T))$
- 7 Computation of $\lim(W(T))$
- 8 Experimentation
- 9 Demo
- 10 Conclusion

Example one

The variable order is $x < y < z$. The regular chain is:

$$\begin{cases} xz - y^2 = 0 \\ y^5 - x^2 = 0 \end{cases}$$

What are the limits of y and z when x approaches 0?

Example one

The variable order is $x < y < z$. The regular chain is:

$$\begin{cases} xz - y^2 = 0 \\ y^5 - x^2 = 0 \end{cases}$$

What are the limits of y and z when x approaches 0?




Figure 1: No limit points at $x = 0$

Example two

The variable order is $x < y < z$. The regular chain is:

$$\begin{cases} xz - y^2 = 0 \\ y^5 - x^3 = 0 \end{cases}$$

What are the limits of y and z when x approaches 0?

Example two

The variable order is $x < y < z$. The regular chain is:

$$\begin{cases} xz - y^2 = 0 \\ y^5 - x^3 = 0 \end{cases}$$

What are the limits of y and z when x approaches 0?




Figure: One limit point at $x = 0$.

How to compute the limit points

The variable order is $x < y < z$.

$$R_1 := \begin{cases} xz - y^2 = 0 \\ y^5 - x^2 = 0 \end{cases}$$

- (1) solve $y^5 - x^2 = 0$, we get $y = x^{\frac{2}{5}}$
- (2) substitute $y = x^{\frac{2}{5}}$ into $xz - y^2 = 0$, we get $xz - x^{\frac{4}{5}} = 0$
- (3) since $x \neq 0$, we have $z = x^{-\frac{1}{5}}$
- (4) so there are no limit points

$$R_2 := \begin{cases} xz - y^2 = 0 \\ y^5 - x^3 = 0 \end{cases}$$

- (1) $y = x^{\frac{3}{5}}$
- (2) $z = x^{\frac{1}{5}}$
- (3) the limit point is $(x = 0, y = 0, z = 0)$

Plan

- 1 The problem
- 2 Motivation
- 3 An introductory example (informal)
- 4 A more advanced example (informal)**
- 5 Limit points and Puiseux expansions of an algebraic curve
- 6 Puiseux expansions of a regular chain and $\lim(W(T))$
- 7 Computation of $\lim(W(T))$
- 8 Experimentation
- 9 Demo
- 10 Conclusion

The problem

- Input: the regular chain R below with $X_1 < X_2 < X_3$

$$R := \begin{cases} r_2 &= (X_1 + 2)X_1X_3^2 + (X_2 + 1)(X_3 + 1) \\ r_1 &= X_1X_2^2 + X_2 + 1 \end{cases}$$

The product of the initials of its polynomials is $h_R := X_1(X_1 + 2)$.

- Output: Limit points of $W(R)$ at $h_R = 0$.

Puiseux series expansions of r_1 at $X_1 = 0$

- The two Puiseux expansions of r_1 at $X_1 = 0$ are:

$$[X_1 = T, X_2 = -1 - T + O(T^2)],$$

$$[X_1 = T, X_2 = -T^{-1} + 1 + T + O(T^2)].$$

- The second expansion cannot result in a limit point while the first one might.

Limit points of $W(R)$ at $X_1 = 0$

- After substituting the first expansion into r_2 , we have:

$$r'_2 = (T + 2)TX_3^2 + (-T + O(T^2))(X_3 + 1)$$

- Now, we compute Puiseux series expansions of r'_2 which are

$$\begin{aligned} [T = T, X_3 = 1 - 1/3 T + O(T^2)], \\ [T = T, X_3 = -1/2 + 1/12 T + O(T^2)]. \end{aligned}$$

- So the regular chains

$$\left\{ \begin{array}{l} X_3 - 1 = 0 \\ X_2 + 1 = 0 \\ X_1 = 0 \end{array} \right\}, \quad \left\{ \begin{array}{l} X_3 + 1/2 = 0 \\ X_2 + 1 = 0 \\ X_1 = 0 \end{array} \right\}$$

give the limit points of $W(R)$ at $X_1 = 0$.

Limit points of $W(R)$ at $X_1 = -2$

- Puiseux series expansions of r_1 at the point $X_1 = -2$:

$$\begin{aligned} [X_1 = T-2, X_2 = 1 + 1/3T + O(T^2)], \\ [X_1 = T-2, X_2 = -1/2 - 1/12T + O(T^2)]. \end{aligned}$$

- After substitution into r_2 , we obtain:



$$\begin{aligned} r'_{12} &= (T-2)TX_3^2 + (2 + 1/3T + O(T^2))(X_3 + 1) \\ r'_{22} &= (T-2)TX_3^2 + (1/2 - 1/12T + O(T^2))(X_3 + 1). \end{aligned}$$

- Puiseux expansions of r'_{12} and r'_{22} at $T = 0$ resulting in limit points:
 - i) for r'_{12} : $[T = T, X_3 = -1 + T + O(T^2)]$
 - ii) for r'_{22} : $[T = T, X_3 = -1 + 4T + O(T^2)]$
- The limit points of $W(R)$ at $X_1 = -2$ are represented by the regular chains $\{X_1 + 2, X_2 - 1, X_3 + 1\}$ and $\{X_1 + 2, X_2 + 1/2, X_3 + 1\}$.

Visualizing the limit points of $W(R)$

The limit points are:

$$\left\{ \begin{array}{l} X_3 - 1 = 0 \\ X_2 + 1 = 0 \\ X_1 = 0 \end{array} \right., \quad \left\{ \begin{array}{l} X_3 + 1/2 = 0 \\ X_2 + 1 = 0 \\ X_1 = 0 \end{array} \right., \quad \left\{ \begin{array}{l} X_3 + 1 = 0 \\ X_2 - 1 = 0 \\ X_1 + 2 = 0 \end{array} \right., \quad \left\{ \begin{array}{l} X_3 + 1 = 0 \\ X_2 + 1/2 = 0 \\ X_1 + 2 = 0 \end{array} \right.$$



Plan


- 1 The problem
- 2 Motivation
- 3 An introductory example (informal)
- 4 A more advanced example (informal)
- 5 Limit points and Puiseux expansions of an algebraic curve**
- 6 Puiseux expansions of a regular chain and $\lim(W(T))$
- 7 Computation of $\lim(W(T))$
- 8 Experimentation
- 9 Demo
- 10 Conclusion

Zariski topology

Zariski closure

- Let $\mathbf{k}$ be an algebraically closed field, like $\mathbb{C}$.
- We denote by $\mathbb{A}^s$ the *affine s -space* over $\mathbf{k}$.
- An *affine algebraic variety* of $\mathbb{A}^s$ is the set of common zeroes of a collection $F \subseteq \mathbf{k}[X_1, \dots, X_s]$ of polynomials.
- The **Zariski topology** on $\mathbb{A}^s$ is the topology whose **closed sets** are the **affine algebraic varieties** of $\mathbb{A}^s$.
- The **Zariski closure** of a subset $W \subseteq \mathbb{A}^s$ is the intersection of all affine algebraic varieties containing W .

The set $\{y = 0, x \neq 0\}$ and its Zariski closure $\{y = 0\}$.



Zariski topology and the Euclidean topology

The relation between the two topologies

- With $\mathbf{k} = \mathbb{C}$, the affine space $\mathbb{A}^s$ is endowed with both topologies.
- The basic open sets of the Euclidean topology are the **open balls**.
- The basic open sets of Zariski topology are the **complements of hypersurfaces**.
- Thus, a Zariski closed (resp. open) set is closed (resp. open) in the Euclidean topology on $\mathbb{A}^s$.
- That is, Zariski topology is **coarser** than the Euclidean topology.

Theorem (The relation between two closures (D. Mumford))

- *Let $V \subseteq \mathbb{A}^s$ be an irreducible affine variety.*
- *Let $U \subseteq V$ be nonempty and open in the Zariski topology induced on V .*

Then, U has the same closure in both topologies. In fact, we have

$$V = \overline{U}^Z = \overline{U}^E.$$

Limit points

Limit points

- Let (X, τ) be a topological space and $S \subseteq X$ be a subset.
- A point $p \in X$ is a **limit point** of S if **every neighborhood** of p contains **at least one point** of S **different from p** itself.
- If X is a **metric space**, the point p is a **limit point** of S if and only if there exists a sequence $(x_n, n \in \mathbb{N})$ of points of $S \setminus \{p\}$ such that $\lim_{n \rightarrow \infty} x_n = p$.
- The **limit points** of S which **do not belong to S** are called **non-trivial**, denoted by $\lim(S)$.

Example

Consider the interval $S := [1, 2) \subset \mathbb{R}$. The point 2 is a non-trivial limit point of S .

Limit points of the quasi-component of a regular chain

Recall Mumford's Theorem

- Let $V \subseteq \mathbb{A}^s$ be an irreducible affine variety.
- Let $U \subseteq V$ be nonempty and open in the Zariski topology induced on V .

Then $V = \overline{U}^Z = \overline{U}^E$.

Corollary

Let R be a regular chain. Recall that $\text{sat}(R) := \langle R \rangle : \text{init}(R)^\infty$ is its saturated ideal and $W(R) = V(R) \setminus V(\text{init}(R))$ is its quasi-component. Then, we have

$$V(\text{sat}(R)) = \overline{W(R)}^Z = \overline{W(R)}^E.$$

- We use $\overline{W(R)}$ to denote this common closure.
- $\lim(W(R)) := \overline{W(R)} \setminus W(R)$ denotes the limit points of $W(R)$.

Field of Puiseux series

- Let T be a symbol.
- $\mathbb{C}[[T]]$: ring of formal power series.
- $\mathbb{C}\langle T \rangle$: ring of convergent power series.
- $\mathbb{C}[[T^*]] = \bigcup_{n=1}^{\infty} \mathbb{C}[[T^{\frac{1}{n}}]]$: ring of formal Puiseux series.
- $\mathbb{C}\langle T^* \rangle = \bigcup_{n=1}^{\infty} \mathbb{C}\langle T^{\frac{1}{n}} \rangle$: ring of convergent Puiseux series.
- $\mathbb{C}((T^*))$: quotient field of $\mathbb{C}[[T^*]]$, or the field of Puiseux series.
- $\mathbb{C}(\langle T^* \rangle)$: quotient field of $\mathbb{C}\langle T^* \rangle$, or the field of convergent Puiseux series.

We have

- $\mathbb{C}[[T]] \subset \mathbb{C}[[T^*]] \subset \mathbb{C}((T^*)); \mathbb{C}\langle T \rangle \subset \mathbb{C}\langle T^* \rangle \subset \mathbb{C}(\langle T^* \rangle)$
- $\mathbb{C}\langle T \rangle \subset \mathbb{C}[[T^*]]; \mathbb{C}\langle T^* \rangle \subset \mathbb{C}[[T^*]]; \mathbb{C}(\langle T^* \rangle) \subset \mathbb{C}((T^*))$

Example

We have $\sum_{i=0}^{\infty} T^i \in \mathbb{C}\langle T \rangle$, $\sum_{i=0}^{\infty} T^{\frac{i}{2}} \in \mathbb{C}\langle T^* \rangle$ and $\sum_{i=-3}^{\infty} T^{\frac{i}{2}} \in \mathbb{C}(\langle T^* \rangle)$.

Theorem (Puisseux)

Both $\mathbb{C}((T^*))$ and $\mathbb{C}(\langle T^* \rangle)$ are algebraically closed fields.

Puisseux expansions

- Let $\mathbf{k} = \mathbb{C}((X^*))$ or $\mathbb{C}(\langle X^* \rangle)$.
- Let $f \in \mathbf{k}[Y]$, where $d := \deg(f, Y) > 0$.
- There exist $\varphi_i \in \mathbf{k}$, $i = 1, \dots, d$, such that

$$\frac{f}{\text{lc}(f, Y)} = (Y - \varphi_1) \cdots (Y - \varphi_d).$$

- We call $\varphi_1, \dots, \varphi_d$ the *Puisseux expansions* of f at the origin.

Example

- $(Y^2 - X) = (Y - X^{\frac{1}{2}})(Y + X^{\frac{1}{2}})$.
- Puiseux expansions of $Y^2 - XY - X$:
 $Y - (X^{\frac{1}{2}} + \frac{1}{2}X + \frac{1}{8}X^{\frac{3}{2}} + O(X^2)), Y - (-X^{\frac{1}{2}} + \frac{1}{2}X - \frac{1}{8}X^{\frac{3}{2}} + O(X^2)).$

Puiseux parametrizations

Let $f \in \mathbb{C}\langle X \rangle[Y]$. A **Puiseux parametrization** of f is a pair $(\psi(T), \varphi(T))$ of elements of $\mathbb{C}\langle T \rangle$ for some new variable T , such that

- $\psi(T) = T^\varsigma$, for some $\varsigma \in \mathbb{N}_{>0}$.
- $f(X = \psi(T), Y = \varphi(T)) = 0$ holds in $\mathbb{C}\langle T \rangle$,
- there is no integer $k > 1$ such that both $\psi(T)$ and $\varphi(T)$ are in $\mathbb{C}\langle T^k \rangle$.

The index ς is the **ramification index** of the parametrization $(T^\varsigma, \varphi(T))$.

Relation to Puiseux expansions

- Let $z_1, \dots, z_\varsigma$ denote the primitive roots of unity of order ς in $\mathbb{C}$. Then $\varphi(z_i X^{1/\varsigma})$, for $i = 1, \dots, \varsigma$, are ς Puiseux expansions of f .
- For a Puiseux expansion φ of f , let c minimum s.t. $\varphi = g(T^{1/c})$ and $g \in \mathbb{C}\langle T \rangle$. Then $(T^c, g(T))$ is a Puiseux parametrization of f .

Example

Puiseux parametrization of $Y^2 - XY - X$:

$$(X = T^2, Y = T + \frac{1}{2} T^2 + \frac{1}{8} T^3 + O(T^4))$$

Plan

- 1 The problem
- 2 Motivation
- 3 An introductory example (informal)
- 4 A more advanced example (informal)
- 5 Limit points and Puiseux expansions of an algebraic curve
- 6 Puiseux expansions of a regular chain and $\lim(W(T))$**
- 7 Computation of $\lim(W(T))$
- 8 Experimentation
- 9 Demo
- 10 Conclusion

Puiseux expansions of a regular chain

Notation

- Let $R := \{r_1(X_1, X_2), \dots, r_{s-1}(X_1, \dots, X_s)\} \subset \mathbb{C}[X_1 < \dots < X_s]$ be a 1-dim regular chain.
- Assume R is strongly normalized, that is, $\text{init}(R) \in \mathbb{C}[X_1]$.
- Let $\mathbf{k} = \mathbb{C}(\langle X_1^* \rangle)$.
- Then R generates a zero-dimensional ideal in $\mathbf{k}[X_2, \dots, X_s]$.
- Let $V^*(R)$ be the zero set of R in $\mathbf{k}^{s-1}$.

Definition

We call *Puiseux expansions* of R the elements of $V^*(R)$.

Remarks

- The *strongly normalized assumption* is only for presentation ease.
- Generically, The 1-dim assumption extends to d -dim $d \leq 2$.
- Higher dimension requires the Jung-Abhyankar theorem.

An example

A regular chain R

$$R := \begin{cases} X_1 X_3^2 + X_2 \\ X_1 X_2^2 + X_2 + X_1 \end{cases}$$

Puiseux expansions of R

$$\begin{cases} X_3 = 1 + O(X_1^2) \\ X_2 = -X_1 + O(X_1^2) \end{cases} \quad \begin{cases} X_3 = -1 + O(X_1^2) \\ X_2 = -X_1 + O(X_1^2) \end{cases}$$

$$\begin{cases} X_3 = X_1^{-1} - \frac{1}{2}X_1 + O(X_1^2) \\ X_2 = -X_1^{-1} + X_1 + O(X_1^2) \end{cases} \quad \begin{cases} X_3 = -X_1^{-1} + \frac{1}{2}X_1 + O(X_1^2) \\ X_2 = -X_1^{-1} + X_1 + O(X_1^2) \end{cases}$$

Relation between $\lim_0(W(R))$ and Puiseux expansions of R

Theorem

For $W \subseteq \mathbb{C}^s$, denote

$$\lim_0(W) := \{x = (x_1, \dots, x_s) \in \mathbb{C}^s \mid x \in \lim(W) \text{ and } x_1 = 0\},$$

and define

$$V_{\geq 0}^*(R) := \{\Phi = (\Phi^1, \dots, \Phi^{s-1}) \in V^*(R) \mid \text{ord}(\Phi^j) \geq 0, j = 1, \dots, s-1\}.$$

Then we have

$$\lim_0(W(R)) = \cup_{\Phi \in V_{\geq 0}^*(R)} \{(X_1 = 0, \Phi(X_1 = 0))\}.$$

$$V_{\geq 0}^*(R) := \left\{ \begin{array}{lcl} X_3 & = & 1 + O(X_1^2) \\ X_2 & = & -X_1 + O(X_1^2) \end{array} \right\} \cup \left\{ \begin{array}{lcl} X_3 & = & -1 + O(X_1^2) \\ X_2 & = & -X_1 + O(X_1^2) \end{array} \right\}$$

Thus the limit ponts are $\lim_0(W(R)) = \{(0, 0, 1), (0, 0, -1)\}$.

Puiseux parametrizations of a regular chain

Idea

- Let $\Phi_i = (\Phi_i^1, \dots, \Phi_i^{s-1}) \in V_{\geq 0}^*(R)$ be a Puiseux expansion, $1 \leq i \leq M := |V_{\geq 0}^*(R)|$. Recall that $\Phi_i^1, \dots, \Phi_i^{s-1} \in \mathbb{C}(\langle X_1^* \rangle)$.
- Φ_i can be associated with a Puiseux parametrization $(X_1 = T^{\varsigma_i}, X_2 = g_i^1(T), \dots, X_s = g_i^{s-1}(T))$ with $g_i^j \in \mathbb{C}\langle T \rangle$.

Details

- Note: Φ_i^j is an expansion of $r_j(X_1, X_2 = \Phi_i^1, \dots, X_j = \Phi_i^{j-1}, X_{j+1})$.
- Let $(T^{\varsigma_{i,j}}, X_j = \varphi_i^j(T))$ be the corresponding Puiseux parametrization of Φ_i^j , where $\varsigma_{i,j}$ is the ramification index of Φ_i^j .
- Let ς_i be the l.c.m. of $\{\varsigma_{i,1}, \dots, \varsigma_{i,s-1}\}$ and $g_i^j := \varphi_i^j(T = T^{\varsigma_i/\varsigma_{i,j}})$.

Definition

$\mathfrak{G}_R := \{(X_1 = T^{\varsigma_i}, X_2 = g_i^1(T), \dots, X_s = g_i^{s-1}(T)), i = 1, \dots, M\}$ is a *system of Puiseux parametrizations* of R .

Relation between $\lim_0(W(R))$ and Puiseux parametrizations of R

Notation (recall)

Let $\mathfrak{G}_R := \{(X_1 = T^{\varsigma_i}, X_2 = g_i^1(T), \dots, X_s = g_i^{s-1}(T)), i = 1, \dots, M\}$
be a **system of Puiseux parametrizations** of R .

Theorem

We have

$$\lim_0(W(R)) = \mathfrak{G}_R(T = 0).$$

Plan

- 1 The problem
- 2 Motivation
- 3 An introductory example (informal)
- 4 A more advanced example (informal)
- 5 Limit points and Puiseux expansions of an algebraic curve
- 6 Puiseux expansions of a regular chain and $\lim(W(T))$
- 7 Computation of $\lim(W(T))$**
- 8 Experimentation
- 9 Demo
- 10 Conclusion

Limit points of a plane curve (without Puiseux parametrizations)

Theorem (Lemaire-MorenoMaza-Pan-Xie 08, $\langle T \rangle \stackrel{?}{=} \text{sat}(T)$)

Let $f \in \mathbb{C}[X][Y]$. Assume that f is primitive in Y . Then $\lim_0(W(f)) = \{(0, y) \mid f(0, y) = 0\}$.

Theorem (R.J. Walker, 50)

Let $f \in \mathbb{C}[X][Y]$. Assume that f is general in Y , that is $f(0, Y) \neq 0$. Then, $\lim_0(W(f)) = \{(0, y) \mid f(0, y) = 0\}$.

Theorem

- Let $f \in \mathbb{C}\langle X \rangle[Y]$.
- Assume that f is general in Y .
- Let $\rho > 0$ be small enough such that f converges in $|X| < \rho$.
- Let $V_\rho(f) := \{(x, y) \mid 0 < |x| < \rho, f(x, y) = 0\}$.

Then, we have $\lim_0(V_\rho(f)) = \{(0, y) \mid f(0, y) = 0\}$.

From algebra to computer: what is the challenge?

Algebra

Let $\mathfrak{G}_R$ be a system of Puiseux parametrizations of R . Recall that we have

$$\lim_0(W(R)) = \mathfrak{G}_R(T = 0).$$

When Walker's theorem applies or when the T is a primitive regular chain, we do not need to compute $\mathfrak{G}_R(T = 0)$. However, those are criteria only!

How to compute $\mathfrak{G}_R$ when the previous criteria do not apply?

- We shall not compute $\mathfrak{G}_R$.
- We need to compute $\mathfrak{G}_R(T = 0)$.
- In fact, we compute a truncation (approximation) of $\mathfrak{G}_R$.

The back-substitution process for computing $\mathfrak{G}_R$

Specifications

Input: $R := \{r_1(X_1, X_2), \dots, r_{s-1}(X_1, \dots, X_s)\}$ a 1-dim strongly normalized regular chain.

Output: $\mathfrak{G}_R$: a system of Puiseux parametrizations of R .

Algorithm

Polynomial	Substitution	Puiseux parametrsation
$r_1(X_1, X_2)$	N/A	$(X_1 = T_1^{s_1}, X_2 = \varphi_1(T_1))$
$r_2(X_1, X_2, X_3)$	$r_2(T_1^{s_1}, \varphi_1(T_1), X_3)$	$(T_1 = T_2^{s_2}, X_3 = \varphi_2(T_2))$
$r_3(X_1, X_2, X_3, X_4)$	$r_3(T_2^{s_1 s_2}, \varphi_1(T_2^{s_2}), \varphi_2(T_2), X_4)$	$(T_2 = T_3^{s_3}, X_4 = \varphi_3(T_3))$

More generally, for $i = 2, \dots, s-1$, we define:

- $f_i := r_i(X_1 = T_1^{s_1}, X_2 = \varphi_1(T_1), \dots, X_i = \varphi_{i-1}(T_{i-1}), X_{i+1}) \in \mathbb{C}\langle T_{i-1} \rangle[X_{i+1}]$,
- $(T_i := T_{i-1}^{s_i}, X_{i+1} := \varphi_i(T_i))$.

New problem: compute Puiseux parametrizations of f_i of given accuracy.

Puiseux parametrizations of $f \in \mathbb{C}\langle X \rangle[Y]$ of finite accuracy

Definition

- Let $f = \sum_{i=0}^{\infty} a_i X^i \in \mathbb{C}[[X]]$.
- For any $\tau \in \mathbb{N}$, let $f^{(\tau)} := \sum_{i=0}^{\tau} a_i X^i$.
- We call $f^{(\tau)}$ the **polynomial part** of f of accuracy $\tau + 1$.

Definition

- Let $f \in \mathbb{C}\langle X \rangle[Y]$, $\deg(f, Y) > 0$.
- Let $\sigma, \tau \in \mathbb{N}_{>0}$ and $g(T) = \sum_{k=0}^{\tau-1} b_k T^k$.
- Let $\{T^{k_1}, \dots, T^{k_m}\}$ be the support of $g(T)$.
- The pair $(T^\sigma, g(T))$ is called a **Puiseux parametrization of f of accuracy τ** if there exists a Puiseux parametrization $(T^\varsigma, \varphi(T))$ of f such that
 - (i) σ divides ς .
 - (ii) $\gcd(\sigma, k_1, \dots, k_m) = 1$.
 - (iii) $g(T^{\varsigma/\sigma})$ is the polynomial part of $\varphi(T)$ of accuracy $(\varsigma/\sigma)(\tau - 1) + 1$.

Computing Puiseux parametrizations of $f \in \mathbb{C}\langle X \rangle[Y]$ of finite accuracy

Theorem

- Let $f = \sum_{i=0}^d \sum_{j=0}^{\infty} a_{i,j} Y^i \in \mathbb{C}\langle X \rangle[Y]$.
- Then we can compute $m \in \mathbb{N}$ such that the Puiseux parametrizations of f of accuracy τ are exactly the Puiseux parametrizations of $\sum_{i=0}^d \sum_{j=0}^{m-1} a_{i,j} Y^i$ of accuracy τ .

Lemma

- Let $f = a_d(X)Y^d + \cdots + a_0(X) \in \mathbb{C}\langle X \rangle[Y]$.
- Let $\delta := \text{ord}(a_d(X))$.
- Then “generically”, we can choose $m = \tau + \delta$.

Recall the back-substitution process for computing $\mathfrak{G}_R$

Algorithm

Polynomial	Substitution	Puiseux parametrisation
$r_1(X_1, X_2)$	N/A	$(X_1 = T_1^{\zeta_1}, X_2 = \varphi_1(T_1))$
$r_2(X_1, X_2, X_3)$	$r_2(T_1^{\zeta_1}, \varphi_1(T_1), X_3)$	$(T_1 = T_2^{\zeta_2}, X_3 = \varphi_2(T_2))$
$r_3(X_1, X_2, X_3, X_4)$	$r_3(T_2^{\zeta_1 \zeta_2}, \varphi_1(T_2^{\zeta_2}), \varphi_2(T_2), X_4)$	$(T_2 = T_3^{\zeta_3}, X_4 = \varphi_3(T_3))$
$\vdots$	$\vdots$	$\vdots$

More generally, for $i = 2, \dots, s-1$, we define:

- $f_i := r_i(X_1 = T_1^{\zeta_1}, X_2 = \varphi_1(T_1), \dots, X_i = \varphi_{i-1}(T_{i-1}), X_{i+1}) \in \mathbb{C}\langle T_{i-1} \rangle[X_{i+1}]$,
- $(T_i := T_{i-1}^{\zeta_i}, X_{i+1} := \varphi_i(T_i))$.

Putting everything together

Let $R := \{r_1(X_1, X_2), \dots, r_{s-1}(X_1, \dots, X_s)\} \subset \mathbb{C}[X_1 < \dots < X_s]$. For $1 \leq i \leq s-1$, let

- $h_i := \text{init}(r_i)$
- $d_i := \deg(r_i, X_{i+1})$
- $\delta_i := \text{ord}(h_i)$.

Theorem

One can compute positive integer numbers $\tau_1, \dots, \tau_{s-1}$ such that, in order to compute $\lim_0(W(R))$, it suffices to compute Puiseux parametrizations of f_i of accuracy τ_i , for $i = 1, \dots, s-1$. Moreover, generically, we can choose τ_i , $i = 1, \dots, s-1$, as follows

- $\tau_{s-1} := 1$
- $\tau_{s-2} := (\prod_{k=1}^{s-2} \varsigma_k) \delta_{s-1} + 1$
- $\tau_i = (\prod_{k=1}^{s-2} \varsigma_k) (\sum_{k=2}^{s-1} \delta_i) + 1$, $i = 1, \dots, s-3$.

Moreover, the indices ς_k can be replaced with d_k , $k = 1, \dots, s-2$.

Plan

- 1 The problem
- 2 Motivation
- 3 An introductory example (informal)
- 4 A more advanced example (informal)
- 5 Limit points and Puiseux expansions of an algebraic curve
- 6 Puiseux expansions of a regular chain and $\lim(W(T))$
- 7 Computation of $\lim(W(T))$
- 8 Experimentation**
- 9 Demo
- 10 Conclusion

Maple packages used: RegularChains and algcurves:-puiseux.

- T : timings of Triangularize
- $\#(T)$: number of regular chains returned by Triangularize
- d-1, d-0: number of one and zero dimensional components
- R : timings spent on removing redundant components
- $\#(R)$: number of irredundant components

Table: Removing redundant components in Kalkbrener decompositions.

Sys	T	$\#(T)$	d-1	d-0	R	$\#(R)$
f-744	14.360	4	1	3	432.567	1
Liu-Lorenz	0.412	3	3	0	216.125	3
MontesS3	0.072	2	2	0	0.064	2
Neural	0.296	5	5	0	1.660	5
Solotareff-4a	0.632	7	7	0	32.362	7
Vermeer	1.172	2	2	0	75.332	2
Wang-1991c	3.084	13	13	0	6.280	13

Plan

- 1 The problem
- 2 Motivation
- 3 An introductory example (informal)
- 4 A more advanced example (informal)
- 5 Limit points and Puiseux expansions of an algebraic curve
- 6 Puiseux expansions of a regular chain and $\lim(W(T))$
- 7 Computation of $\lim(W(T))$
- 8 Experimentation
- 9 Demo**
- 10 Conclusion

Plan

- 1 The problem
- 2 Motivation
- 3 An introductory example (informal)
- 4 A more advanced example (informal)
- 5 Limit points and Puiseux expansions of an algebraic curve
- 6 Puiseux expansions of a regular chain and $\lim(W(T))$
- 7 Computation of $\lim(W(T))$
- 8 Experimentation
- 9 Demo
- 10 Conclusion**

Concluding remarks

- We proposed an algorithm for computing the limit points of the quasi-component of a regular chain in dimension one.
- To this end, we make use of the *Puiseux series expansions* of a regular chain.
- In addition, we have sharp bounds on the degree of truncations that are required to compute *approximate Puiseux series expansions* from which the desired limit points can be obtained.
- Our experimental results show that this is a useful tool for dealing with triangular decompositions of polynomial systems.
- For instance, for testing inclusion between saturated ideals of regular chains in a direct manner (i.e. without computing a basis).
- Computing limit points in higher dimension may require the help of the Abhyankar-Jung theorem. This is work in progress.